


Somerset PCT Tobacco Control Profile

The chart below shows how the indicators for this area compare with the rest of England. The value for each indicator is shown as a circle. The average rate for England is shown by the vertical red line, in the centre of the chart. The range of values for all local areas in England is shown as a grey bar. A red circle means that this area is significantly worse than England for that indicator, Although a green circle shows this area is significantly better than the England average, it may still indicate an important issue in this locality.

The data for these indicators are shown in table one.



Source: Local tobacco control profiles. Produced by the London Health Observatory on behalf of the Public Health Observatories in England. Note: Where there are no values shown, this is because the data are not available or have been suppressed.

Table one: Indicator data for Somerset PCT

Indicator	Value	Regional Average	England Average	England Worst	England Best
Smoking attributable deaths 2007-09	173.6	180.6	216.0	361.5	138.4
Smoking attributable deaths from heart disease 2007-09	23.9	26.4	32.1	59.9	18.7
Smoking attributable deaths from stroke 2007-09	8.2	8.4	10.1	18.4	5.7
Deaths from lung cancer 2007-09	27.2	30.3	38.2	69.4	21.8
Deaths from chronic obstructive pulmonary disease 2007-09	20.5	20.8	26.2	48.7	14.7
Smoking attributable hospital admissions 2009/10	1,168.7	1,285.8	1,417.2	2,508.1	797.3
Cost per capita of smoking attributable admissions, 2009/10	34.8	35.4	37.9	57.7	28.7
Lung cancer registrations 2006-08	35.1	39.3	48.3	87.3	29.7
Oral cancer registrations 2006-08	8.8	8.3	8.9	16.0	5.4
Smoking in pregnancy 2009/10	18.3	14.2	14.0	31.4	4.5
Successful quitters at 4 weeks 2010/11	4,174.7	3,990.4	4,369.8	2,253.3	8,779.1
Successful quitters (CO validated) at 4 weeks 2010/11	3,581.8	3,197.2	3,068.1	281.5	6,172.1
Completeness of NS-SEC recording by Stop Smoking Services 2010/11	73.9	84.2	83.9	50.2	100.0
Prescribed NRT, Varenicline and Bupropion 2009/10	6,100.9	5,688.1	4,847.5	914.7	13,762.9

Source: Local tobacco control profiles. Produced by the London Health Observatory on behalf of the Public Health Observatories in England. Note: Where there are no values shown, this is because the data are not available or have been suppressed.

Supplementary indicators have been provided in a datapack to assist local areas further in their control of tobacco and tobacco related harm. There is no central collection of local level data for some of these indicators. However data may be available locally. This datapack allows users to collect and input their own data for some indicators to make comparisons with national benchmarks. Other indicators in the datapack have been broken down by sex, age and socio-economic group. To download the datapack go to <http://www.lho.org.uk/viewResource.aspx?id=16646>

The tobacco control profiles are part of a series of products produced by the London Health Observatory on behalf of the Public Health Observatories in England that provide local data alongside national comparisons to support local health improvement. For an overview of the health of your local population see the Health Profile for your area at <http://www.healthprofiles.info>

If you would like further information on tobacco and health please go to: http://www.lho.org.uk/LHO_Topics/National_Lead_Areas/NationalSmoking.aspx