

West Somerset

Commissioning Locality profile

Part of the SOMERSET JOINT STRATEGIC NEEDS ASSESSMENT

Practices in Commissioning Locality

Dunster and Porlock Surgeries (Dunster)
Dunster and Porlock Surgeries (Porlock)
Exmoor Medical Centre, Dulverton
Harley House Surgery, Minehead
Irnham Lodge Surgery, Minehead
West Somerset Healthcare, Williton and Watchet



Introduction

This is the seventh Commissioning Locality profile, produced to complement the tenth edition of practice profiles. The document aims to provide an overview of demographic, health and service use data at a Commissioning Locality level.

This document aims to aid Commissioning Localities in the identification of health needs, priorities and potential service interventions that could be commissioned or provided.

A summary of some of the key outcomes is presented first, highlighting the Commissioning Locality's performance compared to the rest of Somerset. This is followed by the Commissioning Locality profile in more detail.

We aim to expand on the intervention summaries highlighting the interventions which can be adopted to improve performance on various outcomes and would particularly welcome feedback on which additional areas would be useful. There are practice level 'ski-slopes' which show the variation of practices within the Commissioning Locality more clearly and again we would welcome feedback on whether you would like to see more (or fewer) of these. Diabetic retinopathy screening data is no longer available to us and there are some changes in definitions and sources of some indicators, otherwise the contents are as in the previous profile. We would be receptive to feedback on any material you feel is less useful and we could consider removing.

The information, together with other data held within the Commissioning Locality can then be used as a rudimentary health needs assessment and could be used to inform decisions within the Commissioning Locality.

Feedback

To give feedback on any aspect of the Commissioning Locality profile please contact the Public Health team.
PublicHealth@somerset.gov.uk

Contents

| | |
|------|--|
| Page | |
| 1 | Presentation of Data |
| 2 | SUMMARY |
| 3 | Brief overview of Commissioning Locality outcomes |
| 5 | POPULATION AND CONTEXT |
| 6 | Demographics - population pyramid |
| 7 | Deprivation scores |
| 8 | People living in deprived areas |
| 9 | Wider determinants of health |
| 11 | DISEASE PREVALENCE |
| 12 | Prevalence |
| 13 | Estimated proportion of people expected to have diabetes who are on register |
| 14 | Estimated proportion of people expected to have dementia who are on register |
| 15 | Estimated proportion of people expected to have COPD who are on register |
| 16 | Respiratory disease profile |
| 17 | Interventions to improve respiratory health |
| 18 | Cardiovascular disease profile |
| 20 | Diabetes profile |
| 21 | Cancer profile |
| 22 | DEATH |
| 23 | Causes of death |
| 24 | Years of life lost |
| 25 | Life expectancy |
| 26 | Standardised Mortality Ratios |
| 28 | Where people die |
| 29 | PREVENTION |
| 30 | NHS Health Checks report |
| 32 | Cancer Screening |
| 35 | Chlamydia screening |
| 36 | Interventions to increase chlamydia screening rate |
| 37 | Seasonal flu vaccinations |
| 38 | Childhood Immunisations |
| 40 | Breastfeeding |
| 41 | Paediatric profile |
| 42 | LIFESTYLE FACTORS |
| 43 | Obesity in schools and in adults |
| 45 | Smoking prevalence |
| 46 | Stop smoking services provision |
| 47 | Smoking evidence based interventions |
| 48 | Alcohol related and drug misuse admissions to hospital and deaths |
| 50 | HOSPITALISATIONS |
| 51 | Standardised admission rates for self-harm and Mental and Behavioural disorders |
| 52 | Standardised admission rates for Falls per 1,000 population aged ≥ 65 and for accidents per 1000 population aged < 18 |
| 53 | Falls and Bone Health |
| 54 | Teenage deliveries in hospital |
| 55 | Standardised emergency admission rate to hospital per 1000 population (ICD grouping) |
| 56 | Emergency admission over 6 years: CHD, CVD, COPD, Asthma, Diabetes |
| 58 | Standardised elective admission rate to hospital per 1000 population (ICD grouping) |
| 59 | Standardised first outpatient attendance rate |
| 60 | PRESCRIBING |
| 61 | MO KTT Prescribing indicators |
| 63 | ASTRO - PUs |

Presentation of data

Data are presented in tables and graphs.

Tables

Tables vary in content, but all contain a comparison of the Commissioning Locality with the other Commissioning Localities and with Somerset. Occasionally there is also a National comparison. The best, median and worst practice values across Somerset are shown where available.

Where possible an indication is made of whether the differences observed are due to chance or are potentially significant:

Green highlighting indicates that the Commissioning Locality value is statistically significantly better than the Somerset average
 Pink highlighting indicates that the Commissioning Locality value is statistically significantly worse than the Somerset average

Occasionally it is not possible to say whether a high value is good or bad; in these instances higher/ lower are used instead of better/worse.

Graphs

The absolute levels of indicators can be on very different scales. In order to show a group of indicators on the same graph the values have been "Normalised" and the plotted value will lie between 0 and 1 - with 0 being the worst and 1 the best (or in some instances lowest and highest)

Normalised score = the difference between the value and the worst value expressed as a percentage of the range of Commissioning Locality values in Somerset

For example if for indicator X the worst value in Somerset is 6 and the best is 11, then the range in Somerset is 5. If the Commissioning Locality has a value of 7 then the normalised score is $(7 - 6)/(11 - 6) = 0.2$

The position of the red diamond indicates where the Commissioning Locality value falls in relation to the other Somerset Commissioning Localities, which are shown as vertical lines.

If the Commissioning Locality markers are not equally spread it means that the distribution of values is not symmetric.

Comments about the Commissioning Locality values will be written in the yellow boxes following the graphs if the Commissioning Locality has an extreme value or if it is significantly higher or lower than the Somerset average.

For some indicators a 'ski slope' of values is shown. This shows the rates, ordered from worst to best, for the county and all of the practices and Commissioning Localities in Somerset. West Somerset and its practices are highlighted. For some other indicators a comparison across the Localities is shown.

Interpretation

When assessing whether a finding needs further investigation it is worth remembering that:


A value can be extreme (eg the "best" or "worst"; "highest" or "lowest") without necessarily being statistically significantly different to the Somerset average.

A statistically significant difference does not necessarily indicate the Commissioning Locality has an extreme value (best or worst; highest or lowest value).

A statistically significant difference might not be a clinically significant difference.

| Example | Position of red marker | Commissioning Locality value is... |
|---------|------------------------|---|
| A | At right hand end | The best (or, for some indicators, the highest) |
| B | Between the ends | An intermediate value |
| C | At left hand end | The worst (or, for some indicators, the lowest) |

Example



West Somerset Commissioning Locality - Summary Public Health Profile 2016

Population & context for health

The area has a much older population profile to the rest of Somerset with proportionally fewer young people and more older people. The Index of Multiple Deprivation for the local area is 23.2 representing higher levels of deprivation compared to the Somerset average of 18.1. More detail on the deprivation data is shown on pp. 7-8 with wider determinants on pp. 9-10.

Disease prevalence

Overviews of respiratory, cardiovascular, diabetes and cancer health issues are shown on pp. 16, 18, 20 and 21 respectively.

Mortality, causes and places of death

The death blobs for mortality and years of life lost on pp.23-4 provide a visual indication of the key causes of death in West Somerset Locality. The proportion of deaths occurring before age 65 years is 10.7% which is similar to the rest of the county, the Somerset rate being 12.0%. Standardised mortality ratios for different conditions and age groups are shown on p.26.

Screening

The commissioning arrangements have changed for the NHS Health Checks programme. Of those eligible, 10.7% received a health check. In the most deprived quintile, 6.4% of the population received a check. A full profile of this data is shown on pp.30-31.

Cervical screening rates for those aged 25-49 years old, at 66.5%, are the lowest / worst in the county of 72.3%. Practice based rates of chlamydia screening of eligible 15-24 year olds was 2.1%, much lower than the Somerset average rate of 3.0%.

Immunisations

Immunisation rates for this year and last are shown on pp.44-5. MMR vaccination rates are above the target for herd immunity of 95%, with 95.0% vaccinated by age 2.

Childhood environment

Local breastfeeding initiation rates are 87.6% compared to 80.7% in Somerset. Continuation rates at 6-8 weeks are 52.3% compared to 49.4% in the county as a whole. The paediatric health profile collates a number of risk factors and outcomes relating to children on p. 41.

Excess weight

According to the underlying model used to assess appropriate childhood weight, only 15% children should be of excess weight. Somerset rates are 22.4% at age 4-5 years and 30.4% at age 10. For this Commissioning

Locality, estimated rates are 22.3% and 30.2% for each age group. The percentage of adults in the area who are obese is 10.5% which is the highest rate in the county.

Smoking

Despite a general reduction in smoking rates in adults in Somerset this remains our greatest public health challenge. The Commissioning Locality has 13.2% of adults over age 16 recorded as smokers compared to a 12.8% Somerset average. However, adult smoking rates have increased slightly in this locality compared to declines in all other localities. In 2015/16, only 35 per 1000 recorded smokers in West Somerset went through cessation services compared to 81 per 1000 in 2013-15. This is a lower than average proportion compared to other Commissioning Localities and this number is dramatically down in absolute terms. Smoking in pregnancy remains a challenge and our county rates are still high compared to the rest of England. More detail on smoking related data is shown on pp. 45-6.

Drugs and Alcohol

Hospital admissions for drug and alcohol related reasons are similar to the Somerset average. Alcohol related admissions: West Somerset 1,833 per 100,000; Somerset 1,894 per 100,000. Drug related admissions: West Somerset 140 per 100,000; Somerset 137 per 100,000.

Hospital admission rates

The hospitalisation section on pp.50-9, contains a wealth of data on reasons why the West Somerset population access hospital services and whether on an emergency or elective basis. To focus on a few indicators, the emergency admission rate for falls in the over 65s is 26 per 1000 as compared to the Somerset average of 31 per 1000. The self-harm admission rate is 206 per 100,000 which is about average for the county.

Medicines Optimisation Key Therapeutic Topics

Performance on the new and revised MOKTT indicators (formerly QIPP indicators) is shown on pp. 61-2. These are designed to promote discussion on the variation, rather than provide targets or influence individual prescribing choices.

Suggested public health areas to prioritise

- Be aware of possible missed cases of dementia.
- Integrate offering chlamydia screening tests within general practice
- Increase cancer screening rates

Please contact the public health team at the council if you would like to discuss further any aspect of your profile or related actions – PublicHealth@Somerset.gov.uk

| Brief overview of Commissioning Locality outcomes | | ENGLAND | Somerset | Commissioning Localities | | | | | | | | |
|---|---|---------|----------|--------------------------|----------------|--------------------------------|-------------|-----------------|----------------|---------|-------------|---------------|
| | | | | Bridgwater | Central Mendip | Crewkerne, Chard and Ilminster | East Mendip | North Sedgemoor | South Somerset | Taunton | West Mendip | West Somerset |
| Population and context | Proportion of population aged 0-14 | 18% | 16% | 18% | 16% | 15% | 17% | 14% | 16% | 17% | 15% | 12% |
| | Proportion of population aged > 75 | 8% | 10% | 9% | 8% | 12% | 8% | 12% | 10% | 10% | 11% | 15% |
| | Index of Multiple Deprivation | 17.4 | 18.1 | 22.4 | 16.6 | 17.3 | 17.8 | 16.2 | 16.9 | 16.8 | 17.2 | 23.2 |
| | Proportion of population living in 10% most deprived neighbourhoods in Somerset* | 24% | 10% | 24% | 8% | 3% | 4% | 8% | 9% | 11% | 4% | 12% |
| | Proportion of population living in 20% most deprived neighbourhoods in Somerset* | 35% | 20% | 36% | 8% | 16% | 25% | 15% | 16% | 15% | 15% | 37% |
| Death | All cause mortality, all ages (SMR compared to Somerset) | | 100% | 98% | 103% | 92% | 99% | 100% | 98% | 109% | 100% | 93% |
| | Proportion of deaths occurring before 65 | 16% | 12% | 14% | 14% | 10% | 12% | 12% | 13% | 11% | 12% | 11% |
| | Proportion of those dying at home (all causes) | 22% | 22% | 24% | 26% | 24% | 26% | 22% | 21% | 19% | 22% | 24% |
| Prevention | Health Checks undertaken as % of eligible (annual target) | | 17% | 11% | 10% | 19% | 36% | 26% | 8% | 16% | 24% | 11% |
| | Health Checks as % of eligible in most deprived population quintile | | 14% | 9% | 5% | 24% | 44% | 21% | 8% | 10% | 28% | 6% |
| | Cervical cancer screening (25-49) | 70.3% | 72.3% | 72.8% | 71.3% | 75.4% | 73.0% | 74.4% | 70.6% | 73.8% | 69.5% | 66.5% |
| | Chlamydia % screened of eligible (15-24 year olds) | | 3.0% | 2.5% | 4.1% | 4.3% | 4.6% | 3.2% | 3.6% | 1.8% | 3.0% | 2.1% |
| | MMR coverage by 2nd birthday | 91.9% | 95.0% | 95.7% | 93.4% | 95.2% | 91.9% | 95.5% | 95.8% | 95.6% | 92.3% | 95.0% |
| | Breastfeeding initiation | | 81% | 75% | 83% | 83% | 82% | 81% | 79% | 82% | 86% | 88% |
| Lifestyle factors | Breastfeeding prevalence at 6-8 weeks | | 49% | 40% | 49% | 50% | 54% | 43% | 49% | 54% | 53% | 52% |
| | Smoking rate adults 16+ | | 12.8% | 17.8% | 14.3% | 13.9% | 12.1% | 12.6% | 11.5% | 11.1% | 10.4% | 13.2% |
| | Smoking ascertainment | | 63% | 67% | 62% | 63% | 60% | 67% | 64% | 61% | 58% | 67% |
| | Smokers going through cessation per 1000 recorded smokers | | 40 | 31 | 40 | 30 | 46 | 38 | 42 | 54 | 40 | 35 |
| | Excess weight in 4-5 year olds | 23% | 22% | 24% | 23% | 26% | 21% | 25% | 23% | 20% | 20% | 22% |
| | Excess weight in 10-11 year olds | 38% | 30% | 32% | 31% | 32% | 29% | 31% | 31% | 28% | 31% | 30% |
| | Obese adults (16+) (rate per 1000 - Standardised to Somerset) | 104 | 88 | 101 | 78 | 94 | 73 | 89 | 86 | 87 | 75 | 105 |
| | Alcohol related admissions (rate per 100,000) | 2,056 | 1,894 | 2,062 | 1,685 | 1,842 | 1,840 | 1,794 | 1,946 | 1,942 | 1,760 | 1,833 |
| Hospital activity | Drug related admissions (rate per 100,000) | | 137 | 157 | 103 | 143 | 119 | 90 | 128 | 161 | 131 | 140 |
| | Teenage deliveries (mother aged <19 at delivery) (rate per 1,000 females aged 15-17) | | 14.7 | 21.5 | 11.1 | 12.7 | 20.3 | 10.5 | 17.6 | 13.5 | 6.4 | 12.6 |
| | Self harm admissions (rate per 100,000) | | 221 | 208 | 199 | 191 | 263 | 135 | 247 | 238 | 218 | 206 |
| | Emergency admissions to hospital for Falls in people aged 65 and over (rate per 1000) | | 31 | 30 | 34 | 28 | 38 | 24 | 34 | 31 | 31 | 26 |

*England data shows what proportion of England's population are in areas with IMD above the Somerset critical value for being in the most deprived 10% or 20%

Red = 'Worst' (most worthy of attention)

Green = 'Best' (least worthy of attention)

| Brief overview of Commissioning Locality outcomes | | ENGLAND | Somerset | Rurality | | Deprivation quintile of practices | | | | | |
|---|---|---------|----------|----------|-------|-----------------------------------|-------|-------|-------|---------------------|---------------|
| | | | | Urban | Rural | 1 most deprived | 2 | 3 | 4 | 5 least deprived | Worse at left |
| | | | | | | | | | | | |
| Population and context | Proportion of population aged 0-14 | 18% | 16% | 17% | 15% | 17% | 16% | 16% | 15% | 16% | |
| | Proportion of population aged > 75 | 8% | 10% | 10% | 11% | 8% | 11% | 10% | 12% | 12% | |
| | Index of Multiple Deprivation | 17.4 | 18.1 | 19.4 | 15.9 | 24.4 | 19.3 | 16.6 | 15.2 | 12.2 | |
| | Proportion of population living in 10% most deprived neighbourhoods in Somerset* | 24% | 10% | 14% | 3% | 28% | 9% | 7% | 0% | 1% | |
| | Proportion of population living in 20% most deprived neighbourhoods in Somerset* | 35% | 20% | 26% | 10% | 44% | 27% | 10% | 7% | 2% | |
| Death | All cause mortality, all ages (SMR compared to Somerset) | | 100% | 107% | 90% | 103% | 107% | 104% | 91% | 90% | |
| | Proportion of deaths occurring before 65 | 16% | 12% | 12% | 11% | 16% | 12% | 12% | 10% | 10% | |
| | Proportion of those dying at home (all causes) | 22% | 22% | 20% | 25% | 23% | 20% | 22% | 24% | 22% | |
| Prevention | Health Checks undertaken as % of eligible (annual target) | | 17% | 19% | 14% | 11% | 20% | 21% | 15% | 14% | |
| | Health Checks as % of eligible in most deprived population quintile | | 14% | 14% | 14% | 9% | 16% | 24% | 8% | 22% | |
| | Cervical cancer screening (25-49) | 70.3% | 72.3% | 71.1% | 74.5% | 70.1% | 70.5% | 73.5% | 73.5% | 75.8% | |
| | Chlamydia % screened of eligible (15-24 year olds) | | 3.0% | 3.1% | 3.0% | 4% | 3% | 3% | 3% | 2% | |
| | MMR coverage by 2nd birthday | 91.9% | 95.0% | 95.1% | 94.8% | 94.9% | 94.2% | 95.4% | 96.0% | 95.0% | |
| | Breastfeeding initiation | | 81% | 80% | 83% | 76% | 80% | 82% | 84% | 84% | |
| | Breastfeeding prevalence at 6-8 weeks | | 49% | 47% | 56% | 43% | 45% | 53% | 56% | 58% | |
| Lifestyle factors | Smoking rate adults 16+ | | 12.8% | 13.8% | 11.2% | 18% | 12% | 12% | 11% | 9% | |
| | Smoking ascertainment | | 63% | 63% | 63% | 64% | 64% | 63% | 61% | 64% | |
| | Smokers going through cessation per 1000 recorded smokers | | 40 | 44 | 32 | 37 | 47 | 46 | 31 | 39 | |
| | Excess weight in 4-5 year olds | 23% | 22% | 22% | 23% | 24% | 22% | 22% | 22% | 21% | |
| | Excess weight in 10-11 year olds | 38% | 30% | 32% | 29% | 33% | 32% | 29% | 29% | 27% | |
| | Obese adults (16+) (rate per 1000 - Standardised to Somerset) | 104 | 88 | 92 | 82 | 104 | 87 | 79 | 74 | 95 | |
| | Alcohol related admissions (rate per 100,000) | 2,056 | 1,894 | 2,010 | 1,712 | 2179 | 1966 | 1851 | 1724 | 1625 | |
| | Drug related admissions (rate per 100,000) | | 137 | 165 | 87 | 199 | 149 | 135 | 96 | 63 | |
| Hospital activity | Teenage deliveries (mother aged <19 at delivery) (rate per 1,000 females aged 15-17) | | 14.7 | 17.9 | 9.6 | 24.1 | 17.1 | 14.4 | 6.5 | 8.2 | |
| | Self harm admissions (rate per 100,000) | | 221 | 250 | 169 | 259 | 249 | 226 | 185 | 142 | |
| | Emergency admissions to hospital for Falls in people aged 65 and over (rate per 1000) | | 31 | 31 | 29 | 31 | 32 | 31 | 29 | 29 | |

*England data shows what proportion of England's population are in areas with IMD above the Somerset critical value for being in the most deprived 10% or 20%

Red = 'Worst' (most worthy of attention)

Green = 'Best' (least worthy of attention)

Population & context for health

The profile contains information on the following aspects of the local population and context for health. These considerations set the scene for health activities within the area and although they don't change much year on year, seeing this background can be helpful in understanding need and planning interventions.

The following information is presented:

- Population pyramid
- Deprivation - as measured using the Index of Multiple Deprivation, see below for an explanation of the seven sub domains which make up this index and the Income Deprivation Affecting Children Index and The Income Deprivation Affecting Older People Index
- People living in highly deprived areas, based on Index of Multiple Deprivation
- Wider context for health – a variety of other indicators such as estimated number of benefits claimants, percentage unemployment and educational attainment within the Commissioning Locality

Index of Multiple Deprivation

The Index of Multiple Deprivation (IMD) combines information from the seven domains to produce an overall relative measure of deprivation. The domains are combined using the following weights:

- Income Deprivation (22.5%)
- Employment Deprivation (22.5%)
- Education, Skills and Training Deprivation (13.5%)
- Health Deprivation and Disability (13.5%)
- Crime (9.3%)
- Barriers to Housing and Services (9.3%)
- Living Environment Deprivation (9.3%)

The weights were derived from consideration of the academic literature on poverty and deprivation, as well as consideration of the levels of robustness of the indicators.

Income Deprivation Domain

The Income Deprivation Domain measures the proportion of the population experiencing deprivation relating to low income. The definition of low income used includes both those people that are out-of-work, and those that are in work but who have low earnings (and who satisfy the respective means tests).

Employment Deprivation Domain

The Employment Deprivation Domain measures the proportion of the working age population in an area involuntarily excluded from the labour market. This includes people who would like to work but are unable to do so due to unemployment, sickness or disability, or caring responsibilities.

Education, Skills and Training Deprivation Domain

The Education, Skills and Training Deprivation Domain measures the lack of attainment and skills in the local population. The indicators fall into two sub-domains: one relating to children and young people and one relating to adult skills.

Health Deprivation and Disability Domain

The Health Deprivation and Disability Domain measures the risk of premature death and the impairment of quality of life through poor physical or mental health. The domain measures morbidity, disability and premature mortality but not aspects of behaviour or environment that may be predictive of future health deprivation.

Crime Domain

The Crime Domain measures the risk of personal and material victimisation at local level.

Barriers to Housing and Services Domain

The Barriers to Housing and Services Domain measures the physical and financial accessibility of housing and local services. The indicators fall into two sub-domains: 'geographical barriers', which relate to the physical proximity of local services, and 'wider barriers' which includes issues relating to access to housing such as affordability and homelessness.

Living Environment Deprivation Domain

The Living Environment Deprivation Domain measures the quality of the local environment. The indicators fall into two sub-domains. The 'indoors' living environment measures the quality of housing; while the 'outdoors' living environment contains measures of air quality and road traffic accidents.

Income Deprivation Affecting Children Index

The Income Deprivation Affecting Children Index (IDACI) measures the proportion of all children aged 0 to 15 living in income deprived families. This is one of two supplementary indices and is a sub-set of the Income Deprivation Domain.

Income Deprivation Affecting Older People Index

The Income Deprivation Affecting Older People Index (IDAOPI) measures the proportion of all those aged 60 or over who experience income deprivation. This is one of two supplementary indices and is a sub-set of the Income Deprivation Domain.


West Somerset

Population at April 2016

33,250


West Somerset

| Age Group | Males | Females |
|-----------|-------|---------|
| 00-04 | 672 | 619 |
| 05-14 | 1,378 | 1,336 |
| 15-24 | 1,594 | 1,553 |
| 25-34 | 1,679 | 1,573 |
| 35-44 | 1,385 | 1,435 |
| 45-54 | 2,142 | 2,229 |
| 55-64 | 2,437 | 2,592 |
| 65-74 | 2,764 | 2,882 |
| 75-84 | 1,544 | 1,848 |
| 85+ | 547 | 1,041 |




Somerset

| Age Group | Males | Females |
|-----------|--------|---------|
| 00-04 | 14,717 | 14,121 |
| 05-14 | 31,431 | 30,024 |
| 15-24 | 31,828 | 29,796 |
| 25-34 | 31,065 | 31,002 |
| 35-44 | 31,179 | 31,952 |
| 45-54 | 41,258 | 41,331 |
| 55-64 | 36,184 | 37,582 |
| 65-74 | 34,633 | 36,370 |
| 75-84 | 18,529 | 21,851 |
| 85+ | 6,620 | 11,869 |



Age/sex Population pyramid. The solid bars represent the population in your Commissioning Locality; solid lines represents Somerset as a whole, dotted lines represents England. Data from Exeter system download of GP registered patients.



The population pyramid shows the percentage of the population at each age group. Conventionally males are shown on the left and females on the right. Each bar represents an age group ordered from youngest at the bottom to oldest at the top.

This data is drawn from the Exeter system download of GP registered patients and the pyramid above shows data for England, Somerset and your Commissioning Locality. The tables to the left show the population numbers in each cohort.

The dotted line represents the population of England as a whole. Due to low death rates in younger years, our population pyramid shows fairly stable proportions in each age cohort (NB the youngest category covers only 5 years). Year on year the pyramid shape can shift due to migration as well as mortality and births.

The solid line for Somerset highlights the greater proportion of people in older age groups.

| Proportion in age range | Commissioning Locality | Somerset | England (mid 2015 projection) | Range of Practice values low / median / high |
|-------------------------|------------------------|----------|-------------------------------|--|
| 0-14 | 12% | 16% | 18% | 9% / 16% / 24% |
| 75+ | 15% | 10% | 8% | 2% / 11% / 19% |
| Female 15-44 | 14% | 16% | 19% | 11% / 16% / 31% |

The Commissioning Locality has the highest value in the county for:
75+

The Commissioning Locality has the lowest value in the county for:
0-14 Female 15-44

Deprivation


Index of Multiple Deprivation (IMD) 2015 and its domains.
 Calculated for Commissioning Locality population in April 2016. All those with a known postcode are included even if they live outside Somerset.

For the IMD score and all of its domains and indices a higher score (and points towards the left of the normalised score graph) indicate a worse situation and more deprivation.


| Score | Commissioning Locality | Somerset | England (median LSOA* values) | Range of Practice values low / median / high |
|---|------------------------|----------|-------------------------------|--|
| IMD score | 23.2 | 18.1 | 17.4 | 9.5 / 16.6 / 26.8 |
| Income score | 0.13 | 0.12 | 0.11 | 0.06 / 0.10 / 0.18 |
| Employment score | 0.12 | 0.10 | 0.10 | 0.05 / 0.09 / 0.15 |
| Health deprivation and disability score | 0.12 | -0.25 | 16.29 | -1.00 / -0.28 / 0.41 |
| Education skills and training score | 27.1 | 20.4 | 0.0 | 6.5 / 19.2 / 40.6 |
| Barriers to housing and services score | 28.9 | 25.2 | 0.0 | 18.4 / 25.2 / 42.6 |
| Crime and disorder score | -0.50 | -0.37 | 20.29 | -1.41 / -0.41 / 0.35 |
| Living environment score | 36.1 | 24.1 | 17.6 | 11.0 / 24.8 / 58.6 |
| Income Deprivation Affecting Children Index | 0.17 | 0.14 | 0.15 | 0.06 / 0.14 / 0.26 |
| Income Deprivation Affecting Older People Index | 0.15 | 0.13 | 0.15 | 0.08 / 0.13 / 0.19 |

*A Lower Super Output Area is a geographical area of about 1500 people.

This bar graph shows where the Commissioning Locality score is on the various deprivation indices compared to the other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. The worst performing Commissioning Locality is the extreme left and best extreme right. Your Commissioning Locality is highlighted with the red diamond. A diamond positioned to the left shows a worse performance.



The Commissioning Locality has the worst value in the county for:
 IMD score Health deprivation and disability score Living environment score



People living in deprived areas

Proportion of Somerset patients registered with the practices in the Commissioning Locality who live in one of the most deprived areas of Somerset and England. Deprivation is measured using the Index of Multiple Deprivation 2015 and calculated for Commissioning Locality populations in April 2016.


Based on the IMD 2015 score, small areas (Lower Super Output Areas - LSOAs) have been rated on their level of deprivation. This has been done for England as a whole and areas grouped into deciles. A grouping has also been done just within Somerset. Somerset is less deprived than England therefore fewer people in Somerset live in a highly deprived area (ie only 8% of the Somerset population lives in the 20% most deprived areas of England and 3% in the 10% most deprived). The population within the Commissioning Locality has been classified by how many live in the 20% and 10% most deprived geographical areas. Differing LSOAs will be in the quintiles and deciles for Somerset and England and may give rise to seemingly contradictory messages in the yellow box below.

Commissioning Localities with a higher proportion of their population living in highly deprived areas will have more challenges with regards to health interventions.

All patients with a known LSOA are included below (if they are living outside Somerset their IMD score is compared with the Somerset deprivation quintiles for the first two indicators)

| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|--|------------------------|----------|---------|--|
| Proportion of population in 20% most deprived areas (Somerset) | 37% | 20% | 35% | 0% / 12% / 77% |
| Proportion of population in 10% most deprived areas (Somerset) | 12% | 10% | 24% | 0% / 2% / 40% |
| Proportion of population in 20% most deprived areas (England) | 0% | 8% | 20% | 0% / 1% / 37% |
| Proportion of population in 10% most deprived areas (England) | 0% | 3% | 10% | 0% / 0% / 17% |

This bar graph shows the proportion of the people in the Commissioning Locality who live in the areas rated in the 10% and 20% most deprived areas of England and/or Somerset. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted with the red diamond. The worst performing Commissioning Locality score is the extreme left and indicates a higher proportion of the population living in very deprived areas and the best is on the extreme right indicating a low proportion living in very deprived areas.




The Commissioning Locality has the worst value in the county for:

Proportion of population in 20% most deprived areas (Somerset)

The Commissioning Locality has the best value in the county for:

Proportion of population in 20% most deprived areas (England)



Wider determinants of health


Population weighted estimates of indicators available at Lower Super Output level except for the final two indicators which are by Commissioning Locality geographical area

This set of data gives more clue as to the specific issues which affect the Commissioning Locality population such as health status, proportion unemployed and benefits claimants. The data source and time period for each indicator is given in the table below and uses population weighted estimates of indicators available at Lower Super Output (LSOA) level, a small grouping of about 1500 people. England figures are not available for all indicators and some may not be for exactly the same time period.

To calculate the Commissioning Locality value, each registered patient is assumed to have the same pattern of experience as the whole LSOA in which they live, thus they are not exact measures but based on averages. As the populations of the localities have changed the value shown for indicators from the Census will have changed since the last profile was published. The indicators are summarised graphically on the following page.

| | Commissioning Locality | Somerset | England | Range of Commissioning Locality values |
|--|------------------------|----------|---------|--|
| % of Residents that are not White British Census 2011 | 4% | 5% | 20% | 4% to 7% |
| % of People whose day-to-day activities are limited a lot Census 2011 | 11% | 8% | 9% | 7% to 11% |
| % of People whose health is bad or very bad Census 2011 | 7% | 5% | 6% | 4% to 7% |
| % of People who provide unpaid care Census 2011 | 13% | 11% | 10% | 10% to 13% |
| % of People aged 16 or over with no qualifications Census 2011 | 25% | 23% | 23% | 21% to 26% |
| % of Households that are socially rented Census 2011 | 15% | 14% | 18% | 9% to 16% |
| % of Households that are lone parent households Census 2011 | 7% | 9% | 11% | 7% to 10% |
| % of Households with No Access to car/van Census 2011 | 19% | 16% | 26% | 13% to 19% |
| % of Residents aged 16-74 who are Unemployed Census 2011 | 2.8% | 3.0% | 4.4% | 2.6% to 4.1% |
| Foundation Stage Profile (FSP) % students NOT achieving a good level of development Somerset County Council 2015 | 50% | 33% | n/a | 28% to 50% |
| % of pupils with SEN Somerset County Council 2015 | 15% | 15% | n/a | 12% to 17% |
| Fixed exclusions per 1000 pupils Somerset County Council 2015 | 41 | 46 | n/a | 31 to 52 |
| % of children not achieving 5 A*-C GCSEs including Maths and English Somerset County Council 2015 | 50% | 41% | n/a | 36% to 50% |
| Community Services recipients aged 75+ (rate per 1000 aged 75+) SCC Adult social care - Mar16 | 24 | 33 | n/a | 24 to 51 |
| Home Care recipients aged 65+ (rate per 1000 aged 65+) SCC Adult social care - Mar16 | 14 | 18 | n/a | 14 to 24 |
| Applications for housing on Homefinder SCC % of census households - Mar16 | 4% | 3% | n/a | 3% to 5% |
| Criminal Damage (rate per 100,000 population) Home Office 2015/16 | 523 | 712 | n/a | 523 to 959 |
| Violence and Sexual offences (rate per 1000 census households) Home Office 2015/16 | 29.0 | 36.9 | n/a | 28.2 to 55.0 |
| % of Children in low-income families Child Poverty Unit 2014 | 17% | 15% | 20% | 13% to 18% |
| % of Households in Fuel Poverty DECC 2014 | 14% | 13% | 11% | 11% to 14% |
| Attendance Allowance claimants (% of population aged 65+) NOMIS Feb16 | 14% | 13% | 14% | 11% to 14% |
| Job Seekers Allowance (% of working population) NOMIS Feb16 | 0.5% | 0.8% | 1.5% | 0.5% to 1.1% |
| Income Support or Pension Credit claimants (% of total population) NOMIS Feb16 | 5% | 4% | 4% | 3% to 5% |
| Pension Credit claimants (% of population aged 65+) NOMIS Feb16 | 15% | 14% | 17% | 12% to 17% |
| Council tax bands A to C (% of households) 2016 | 62% | 66% | 66% | 59% to 75% |
| Children 0-17 currently in care known to be living or placed in Somerset (rate per 10,000) Somerset County Council as at Jun16 | 33 | 35 | 60 | 19 to 44 |
| Children 0-17 subject to a Child Protection Plan known to be living or placed in Somerset (rate per 10,000) Somerset County Council as at Jun16 | 22 | 30 | 43 | 7 to 61 |

This bar graph shows where the Commissioning Locality score is on the various indicators compared to the other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted with the red diamond. The worst performing Commissioning Locality score is the extreme left and best extreme right.



The Commissioning Locality has the worst/highest value in the county for:

Activities limited a lot Health bad or very bad Unpaid carers No Access to car/van Foundation Stage Profile good level NOT achieved Children NOT achieving 5 good GCSEs Income Support or Pension Credit (all ages)

The Commissioning Locality has the best/lowest value in the county for:

Lone parent households Community Services recipients aged 75+ per 1000 Home Care recipients aged 65+ per 1000 Criminal Damage per 1000 households Job Seekers Allowance

Disease prevalence

The profile contains information on the following aspects of the local population and their level of disease. Dependent on the data this information can give an indication of need within the population and success of case finding programmes.

The following information is presented:

- Prevalence numbers in the Commissioning Locality from practice clinical systems compared to expected numbers if the Commissioning Locality prevalence mirrored the average recorded rates over Somerset (adjusted for age and sex differences)
- Change over time in observed numbers of people on the disease registers compared to modelled expected numbers: diabetes, dementia & COPD
- Disease profiles, an overview of factors contributing to disease levels, see below

Presentation of actual disease prevalence is based on MIQUEST and can inform commissioning plans to cover the number of people with the condition. However when calculating level of disease it is useful to know how this compares to average levels which might be expected to expose possible undiagnosed cases and/or particular health needs in the area. There are two methods of calculating expected prevalence of disease used in this profile.

The crude prevalence rate (such as appeared in QOF rates) is calculated by comparing the Commissioning Locality and Somerset values of the number recorded with the condition as a proportion of the total population. The first method shown on the next page is to improve upon this estimate by adjustments to allow for the extent to which the Commissioning Locality has a different age and sex profile to Somerset as a whole. This is important where the disease has a clear variation by age or gender, for example in dementia. Using this approach, some Commissioning Localities will always be under diagnosing whereas others will always be over diagnosing relative to the Somerset average.

The second method of estimating prevalence comes from assumptions about the true prevalence of disease. Using this second approach it would be possible for the whole of Somerset to be over or under diagnosing. Levels of prevalence below expected levels may indicate undiagnosed cases or it could indicate an area in which the Commissioning Locality has better than expected population health.

A comparison of the recorded prevalence to the modelled prevalence in the Commissioning Locality is often used as an estimate of diagnosis completeness. Here we have focused on three conditions - diabetes, dementia and COPD - and shown how the estimated diagnosis levels have changed over time.

It is important to note that the assumptions made by the model are critical and revisions to the model may affect quite dramatically the expected numbers and thus diagnosis levels.

Disease profiles were introduced to provide a useful collation of a range of data on a particular area. The aim is to enable a more holistic picture to be built up of the most appropriate actions along the disease pathway to improve health.

The profiles are designed to provide information on:

- Prevalence of relevant conditions
- Mortality rates
- Preventative intervention
- Risk factors
- Admissions to hospital
- QOF On-going management indicators

Profiles have been produced on:

- Respiratory disease
- Cardiovascular disease
- Diabetes
- Cancer

We would particularly welcome feedback on these profiles, whether they are useful, what you like, aspects which may be less clear and suggestions for additional indicators to include or diseases to cover. Please email PublicHealth@somerset.gov.uk

**Prevalence rate
(rate per 1000 population)**

Age/sex standardised prevalence rates using Somerset wide age/sex specific rates from MIQUEST (using the Quality Outcome Framework definitions) and Exeter system population downloads .


This data shows a comparison of the level of recorded disease in the Commissioning Locality compared to the level in the totality of Somerset practices. The numbers of patients with recorded disease at the practice level has been derived from the Quality Outcome Framework submissions, reported annually through the Information Centre website. This is then converted to a rate per 1000 of the population based on Exeter system population size and composition for each practice which has been retrieved by MIQUEST. MIQUEST is a locally used interrogation tool that retrieves data from practice clinical systems.

The level of disease /condition expected in the Commissioning Locality has been calculated by taking the total Somerset recorded numbers and then partialling this out to the number expected in the Commissioning Locality, making allowances for differences in the age and sex profile of the Commissioning Locality compared to Somerset as a whole. This adjustment is particularly important where conditions are more common in specific age groups or by gender.

2016

| Condition | Observed in Locality | Expected in Locality | Commissioning Locality rate | Somerset rate | England rate (2015) | Range of Practice values low / median / high |
|------------------------------|----------------------|----------------------|-----------------------------|---------------|---------------------|--|
| Asthma | 2,112 | 2,203 | 61.8 | 64.5 | 63.8 | 46 / 66 / 77 |
| Atrial Fibrillation | 1,191 | 1,126 | 26.1 | 24.6 | 24.4 | 16 / 25 / 31 |
| Chronic Kidney Disease (18+) | 1,671 | 1,760 | 44.3 | 46.7 | 57.6 | 25 / 44 / 80 |
| COPD | 881 | 937 | 20.2 | 21.5 | 25.2 | 10 / 21 / 38 |
| Coronary Heart Disease | 1,745 | 1,680 | 39.0 | 37.6 | 44.7 | 22 / 37 / 55 |
| Dementia | 303 | 456 | 6.3 | 9.5 | 11.4 | 4 / 9 / 18 |
| Diabetes (17+) | 2,091 | 2,266 | 60.8 | 65.9 | 82.6 | 48 / 65 / 95 |
| Epilepsy (18+) | 279 | 253 | 9.6 | 8.7 | 8.7 | 4 / 8 / 18 |
| Heart Failure | 370 | 407 | 8.0 | 8.8 | 11.0 | 5 / 9 / 19 |
| Hypertension | 6,479 | 6,944 | 150 | 161 | 184.8 | 130 / 161 / 228 |
| Learning Difficulties (18+) | 189 | 133 | 7.2 | 5.1 | 5.5 | 0 / 4 / 19 |
| Mental Health | 255 | 253 | 7.3 | 7.2 | 9.8 | 3 / 6 / 17 |
| Obesity (16+) | 3,307 | 2,778 | 105 | 88 | 103.7 | 33 / 81 / 235 |
| Osteoporosis (50+) | 52 | 64 | 2 | 3 | 3.6 | 0 / 2 / 15 |
| Palliative Care | 159 | 123 | 4.3 | 3.3 | 5.9 | 0 / 3 / 14 |
| Rheumatoid Arthritis (16+) | 260 | 299 | 7.5 | 8.6 | 9.2 | 4 / 8 / 12 |
| Stroke | 1,378 | 1,219 | 30.9 | 27.3 | 24.1 | 20 / 27 / 34 |
| Thyroid | 1,642 | 1,603 | 40.6 | 39.6 | n/a | 22 / 40 / 51 |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their observed and expected numbers. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show more cases recorded than expected compared to other Commissioning Localities and values to the right show much fewer cases recorded than expected. Values to the left may indicate an increased burden of disease whereas values to the right may indicate under-diagnosis and where case finding efforts may need to be increased.



Significantly worse (higher number than expected compared to the county average) for:

Learning Difficulties (18+) Obesity (16+) Palliative Care Stroke

The Commissioning Locality has the worst value (highest number compared to the expected based on county average) in the county for:

Atrial Fibrillation Learning Difficulties (18+) Obesity (16+) Stroke

Significantly better (lower number than expected compared to the county average) for:

Dementia Diabetes (17+) Hypertension

The Commissioning Locality has the best value (lowest number compared to the expected based on county average) in the county for:

Dementia Heart Failure Hypertension Rheumatoid Arthritis (16+)

Estimated proportion of people expected to have Diabetes who are on register

Register data compared with modelled Type1 and Type 2 prevalence. Data based on Yorkshire and Humberside Public Health Observatory model.

This data shows a comparison of the actual level of disease with the "true" level of the condition estimated to be in the population. The numbers of patients with recorded disease at the practice level has been derived from the Quality Outcome Framework submissions, reported annually through the Information Centre website. Note that these numbers vary slightly from those from local MIQUEST queries used in the Prevalence section and Cardiovascular and Diabetes profiles.



The expected values are based on the Yorkshire and Humberside Public Health Observatory Model

An observed value below that of the expected value may indicate there are undiagnosed patients with that condition within the Commissioning Locality population. Conversely a higher number in the observed column may also warrant further investigation and may indicate areas of high prevalence or possibly over diagnosis.

The ratio can be greater than 100% as it is the comparison of the actual number on the register compared to a modelled expected number.

| | | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|------|------------------------------------|------------------------|----------|-----------|--|
| 2015 | Observed number on register | 2,047 | 29,656 | 2,703,044 | |
| | Expected number on register | 3,306 | 45,322 | 3,321,750 | |
| | Proportion of expected on register | 61.9% | 65.4% | 81.4% | 47% / 65% / 88% |
| 2014 | Observed number on register | 1,997 | 28,516 | 2,703,044 | |
| | Expected number on register | 2,642 | 36,265 | 3,321,750 | |
| | Proportion of expected on register | 75.6% | 78.6% | 81.4% | 56% / 78% / 110% |
| 2013 | Observed number on register | 1,904 | 27,046 | 2,703,044 | |
| | Expected number on register | 2,591 | 35,248 | 3,321,750 | |
| | Proportion of expected on register | 73.5% | 76.7% | 81.4% | 53% / 76% / 109% |
| 2012 | Observed number on register | 1,792 | 25,624 | 2,566,436 | |
| | Expected number on register | 2,578 | 34,845 | 3,245,432 | |
| | Proportion of expected on register | 69.5% | 73.5% | 79.1% | 51% / 71% / 118% |
| 2011 | Observed number on register | 1,717 | 24,405 | 2,455,937 | |
| | Expected number on register | 2,527 | 33,771 | 3,166,556 | |
| | Proportion of expected on register | 67.9% | 72.3% | 77.6% | 51% / 71% / 118% |

The bar chart shows how Commissioning Locality performance compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show a lower ratio of observed to expected cases, in general indicating under-diagnosis compared to the model. Values to the right show a higher ratio of observed to expected cases although even so this does not exclude the possibility of some under reporting.



Estimated proportion of people expected to have Dementia who are on register

Register data compared with modelled prevalence as reported on the Primary Care Web tool www.primarycare.nhs.uk, a different source to that used in previous profiles

This data shows a comparison of the actual level of disease with the "true" level of the condition estimated to be in the population. The numbers of patients with recorded disease at the practice level has been derived from the Quality Outcome Framework submissions, reported annually through the Information Centre website. Note that these numbers vary slightly from those from local MIQUEST queries used in the Prevalence section.


The expected values are based on the NHS Information Centre model as found on the NHS comparators website.

An observed value below that of the expected value may indicate there are undiagnosed patients with that condition within the Commissioning Locality population. Conversely a higher number in the observed column may also warrant further investigation and may indicate areas of high prevalence or possibly over diagnosis.

The ratio can be greater than 100% as it is the comparison of the actual number on the register compared to a modelled expected number.

| | | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|------|------------------------------------|------------------------|----------|---------|--|
| 2015 | Observed number on register | 309 | 5,284 | 415,964 | |
| | Expected number on register | 690 | 8,938 | 684,371 | |
| | Proportion of expected on register | 44.8% | 59.1% | 60.8% | 24% / 56% / 122% |
| 2014 | Observed number on register | 252 | 4,494 | 348,973 | |
| | Expected number on register | 708 | 8,716 | 674,309 | |
| | Proportion of expected on register | 35.6% | 51.6% | 51.8% | 16% / 52% / 91% |
| 2013 | Observed number on register | 248 | 4,178 | 318,669 | |
| | Expected number on register | 752 | 8,472 | 662,645 | |
| | Proportion of expected on register | 33.0% | 49.3% | 48.1% | 17% / 48% / 96% |
| 2012 | Observed number on register | 225 | 3,681 | 293,729 | |
| | Expected number on register | 742 | 8,342 | 656,240 | |
| | Proportion of expected on register | 30.3% | 44.1% | 44.8% | 0% / 40% / 88% |
| 2011 | Observed number on register | 195 | 3,211 | 266,547 | |
| | Expected number on register | 672 | 7,497 | 602,234 | |
| | Proportion of expected on register | 29.0% | 42.8% | 44.3% | 0% / 40% / 88% |

The bar chart shows how Commissioning Locality performance compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show a lower ratio of observed to expected cases, in general indicating under-diagnosis compared to the model. Values to the right show a higher ratio of observed to expected cases although even so this does not exclude the possibility of some under reporting.



The Commissioning Locality has the lowest value in the county for:
 2015 2014 2013 2012 2011

Estimated proportion of people expected to have COPD who are on register

Register data compared with modelled prevalence using models from the NHS Comparators website.

This data shows a comparison of the actual level of disease with the "true" level of the condition estimated to be in the population. The numbers of patients with recorded disease at the practice level has been derived from the Quality Outcome Framework submissions, reported annually through the Information Centre website . Note that these numbers vary slightly from those from local MIQUEST queries used in the Prevalence section and in the Respiratory profile.


The expected values are based on NHS Information Centre models as found on the NHS comparators website, except for diabetes which uses the Yorkshire and Humber Public Health Observatory Model.

An observed value below that of the expected value may indicate there are undiagnosed patients with that condition within the Commissioning Locality population. Conversely a higher number in the observed column may also warrant further investigation and may indicate areas of high prevalence or possibly over diagnosis.

The ratio can be greater than 100% as it is the comparison of the actual number on the register compared to a modelled expected number.

| | | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|------|------------------------------------|------------------------|----------|-----------|--|
| 2015 | Observed number on register | 832 | 11,351 | 1,034,578 | |
| | Expected number on register | 847 | 11,487 | 955,221 | |
| | Proportion of expected on register | 98.2% | 98.8% | 108.3% | 49% / 96% / 163% |
| 2014 | Observed number on register | 790 | 11,029 | 1,004,920 | |
| | Expected number on register | 833 | 11,284 | 940,097 | |
| | Proportion of expected on register | 94.8% | 97.7% | 106.9% | 42% / 93% / 168% |
| 2013 | Observed number on register | 771 | 10,507 | 974,999 | |
| | Expected number on register | 819 | 10,994 | 926,224 | |
| | Proportion of expected on register | 94.1% | 95.6% | 105.3% | 50% / 93% / 157% |
| 2012 | Observed number on register | 717 | 9,924 | 938,511 | |
| | Expected number on register | 805 | 10,751 | 907,873 | |
| | Proportion of expected on register | 89.0% | 92.3% | 103.4% | 52% / 86% / 167% |
| 2011 | Observed number on register | 645 | 9,487 | 898,989 | |
| | Expected number on register | 791 | 10,490 | 888,795 | |
| | Proportion of expected on register | 81.6% | 90.4% | 101.1% | 52% / 86% / 167% |

The bar chart shows how Commissioning Locality performance compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show a lower ratio of observed to expected cases, in general indicating under-diagnosis compared to the model. Values to the right show a higher ratio of observed to expected cases although even so this does not exclude the possibility of some under reporting.




Respiratory data



West Somerset

| | |
|--|---|
| | Significantly better than Somerset average |
| | Not significantly different to Somerset average |
| | Significantly worse than Somerset average |
| | Difference not assessed |



| | |
|--|----------|
| | Improved |
| | Worsened |

| Significance | Indicator | West Somerset number | West Somerset value | Somerset average | Worst Locality value | Locality range ←Worse Better→ | Best Locality value | West Somerset value 2015 | Direction of change |
|--|--|----------------------|---------------------|------------------|----------------------|----------------------------------|---------------------|--------------------------|---------------------|
| Prevalence | 1 : Asthma | 2,112 | 61.8 | 64.5 | 70.9 | | 61.6 | 62.9 | |
| | 2 : COPD | 881 | 20.2 | 21.5 | 26.2 | | 18.6 | 19.5 | |
| | 3 : COPD estimated proportion of cases on register | 832 | 98% | 99% | 85% | | 116% | 95% | |
| Mortality | 4 : All respiratory disease | 223 | 77% | 100% | 123% | | 77% | 76% | |
| | 5 : COPD | 81 | 71% | 100% | 118% | | 71% | 70% | |
| Flu vaccination | 6 : 65 and over | 7,022 | 65% | 70% | 65% | | 74% | 70% | |
| | 7 : All children aged 2-4 | 340 | 41% | 42% | 32% | | 47% | 42% | |
| | 8 : At risk 6 months to <65 years | 1,494 | 42% | 46% | 42% | | 48% | 48% | |
| | 9 : Pregnant women | 120 | 38% | 43% | 33% | | 55% | 36% | |
| Smoking | 10 : Carers | 85 | 26% | 37% | 25% | | 44% | 31% | |
| | 11 : Current smokers (aged 16 and over) | | 13% | 13% | 18% | | 10% | 14% | |
| | 12 : 4 week smoking quit rate (all ages) | | 58% | 49% | 40% | | 60% | 43% | |
| | 4 week smoking CO confirmed quit rate (all ages) | | 40% | 37% | 29% | | 53% | | New indicator |
| | 14 : 4 week smoking quit rate (45-64) | | 42% | 45% | 30% | | 66% | 45% | |
| | 15 : Smokers going through cessation per 1000 recorded smokers | | 35.1 | 40.2 | 30.3 | | 54.3 | 80.6 | |
| Emergency admissions | 16 : Emergency admissions for Respiratory diseases (all ages) | 525 | 13.7 | 13.6 | 15.0 | | 12.0 | 9.5 | |
| | 17 : Emergency admissions for Respiratory diseases (<18) | 422 | 13.9 | 13.4 | 15.4 | | 10.9 | 13.3 | |
| | 18 : Emergency admissions for COPD (all ages) | 253 | 1.9 | 2.2 | 2.6 | | 1.6 | 1.6 | |
| | 19 : Emergency admissions for Asthma (all ages) | 67 | 0.7 | 0.8 | 0.9 | | 0.6 | 0.7 | |
| | 20 : Emergency admissions for Asthma (<18) | 41 | 1.3 | 1.5 | 1.9 | | 1.3 | 1.4 | |
| QOF 2015 ongoing management indicators | 21 : AST004. Patients with asthma aged 14 - 19 years with record of smoking status | 253 | 84% | 68% | 57% | | 88% | 75% | |
| | 22 : AST003. Had a review | 67 | 71% | 78% | 50% | | 77% | 67% | |
| | 23 : COPD004. Record of FEV1 | 41 | 135% | 155% | 65% | | 89% | 81% | |
| | 24 : COPD003. Review including MRC dyspnoea score | 83 | 84% | 68% | 64% | | 94% | 84% | |
| | 25 : COPD005. MRC dyspnoea grade ≥3, with a record of oxygen saturation value | 1,360 | 72% | 59% | 74% | | 97% | 92% | |
| | 26 : COPD007. Had influenza immunisation | 627 | 84% | 74% | 87% | | 99% | 94% | |

- Indicator Notes
- 1-2 MIQUEST (QOF) indirectly standardised prevalence rate within Somerset 2016
 - 3 Modelled true prevalence using Doncaster models compared to QOF reported cases 2015
 - 4-5 Indirectly standardised mortality ratio (compared to Somerset) : ONS : 2011-15
 - 6-10 Uptake of Flu vaccination : PHE : Winter of 2015/16
 - 11-15 Smoking prevalence from MIQUEST query March 2016 and SmokefreeLife Somerset service data April 2015 to March 2016
 - 16-20 Indirectly standardised admission to hospital rate per 100,000 : Secondary Uses Service (SUS) : 2015/16 for Respiratory (all ages) and Asthma (all ages), 2013/14 - 2015/16 for COPD (all ages) and Asthma (<18).
Respiratory diseases ICD10 codes: Chapter J
COPD ICD10 codes: J40-J44
Asthma ICD10 code J45, J46
 - 17-22 QOF ongoing management indicators : 2015

AST004: The percentage of patients with asthma aged 14 or over and who have not attained the age of 20, on the register, in whom there is a record of smoking status in the preceding 12 months
 AST003: The percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months that includes an assessment of asthma control using the 3 RCP questions, NICE 2011 menu ID: NM23
 COPD004: The percentage of patients with COPD with a record of FEV1 in the preceding 12 months
 COPD003: The percentage of patients with COPD who have had a review, undertaken by a healthcare professional, including an assessment of breathlessness using the Medical Research Council dyspnoea scale in the preceding 12 months
 COPD005: The percentage of patients with COPD and Medical Research Council dyspnoea grade ≥3 at any time in the preceding 12 months, with a record of oxygen saturation value within the preceding 12 months, NICE 2012 menu ID: NM63
 COPD007: The percentage of patients with COPD who have had influenza immunisation in the preceding 1 August to 31 March

Interventions to improve respiratory health

Respiratory disease is one of the key contributing factors to reduced life expectancy in Somerset and there are variations and inequalities in the experience of respiratory illness across the county with a clear relationship between deprivation and poor respiratory health.

Priorities to be considered for respiratory services:

- Increase number of patients referred for stop smoking support to Smokefree Somerset <http://www.smokefreeifomeset.co.uk/> and encouragement to patients not ready to quit to have a smokefree home, especially parents of young children.
- For patients not ready to quit signpost to the NHS Smokefree website <https://www.nhs.uk/smokefree/>. Patients interested in vaping instead of smoking should not be discouraged as the risks are substantially lower <https://www.gov.uk/government/news/e-cigarettes-around-95-less-harmful-than-tobacco-estimates-landmark-review>.

Respiratory disease has a number of risk factors that can be minimised and approaches that can add to the effective management of the disease. Overall aims of respiratory services should look to:

- Increase the number of people with long term conditions living independently and in control of their condition e.g. COPD through managed self-care. People with COPD/asthma should be reviewed regularly by a nurse/doctor with appropriate training in respiratory disease management. Review should incorporate a written action plan¹. Further review(s) can take place opportunistically:
 - An acute consultation offers opportunity to determine what action the patient has already taken to deal with the exacerbation. Their self-management strategy may be reinforced or refined and the need for consolidation at a routine follow up considered.
 - A consultation for an upper respiratory tract infection is an opportunity to rehearse self-management in the event of their condition deteriorating.
- Encourage compliance with prescribed medications, with information and support on use and effectiveness, for example;
 - Prescribe inhalers only after patients have received training in the use of the device and have demonstrated satisfactory technique²
 - If the patient is unable to use a device satisfactorily an alternative should be found.
 - The patient should have their ability to use an inhaler device assessed by a competent healthcare professional.
 - The medication needs to be titrated against clinical response to ensure optimum efficacy.
 - Reassess inhaler technique as part of structured clinical review
- Increase rates of immunisation against seasonal flu and pneumonia amongst eligible groups; Research³ has shown the following approaches effective:
 - Having a lead member of staff to oversee the vaccination programme;
 - Ordering sufficient vaccine for 75% uptake among eligible groups;
 - Tailor call-recall process: Personal invitations (particularly effective for the over 65s) and repeats/reminders (particularly effective for under 65s);
 - Using in house IT search to identify eligible patients;
 - Community midwives administering vaccine to pregnant women;
 - Production of report reviewing annual vaccination rates, including detail of rationale to end each annual campaign.
- Increase awareness in the population of signs and symptoms of respiratory disease and when to seek medical advice, seeking to minimise emergency admissions, particularly amongst children and the most deprived groups.
 - Case detection studies have used symptom questionnaires to screen for asthma in school-age children^{2,3,4,5}. It is recommended¹ to focus the initial assessment in children suspected of having asthma on:
 - Presence of key features in the history and examination
 - Careful consideration of alternative diagnoses

1] Galant SP, Crawford LJ, Morpew T, Jones CA, Bassin S. (2004) Predictive value of a cross-cultural asthma case-detection tool in an elementary school population. *Pediatrics* 114(3) pp: 307-16.

2] Healthcare Improvement Scotland (2012) British Guidelines on the Management of Asthma. Scottish Intercollegiate Guidelines Network. Available from <http://www.brit-thoracic.org.uk/Portals/0/Guidelines/AsthmaGuidelines/sign101%20Jan%202012.pdf>

3] Dexter, L. J., Teare, M. D., Dexter, M., Siriwardena, A. N. and Read, R. C. (2012) Strategies to increase influenza vaccination rates: outcomes of a nationwide cross sectional survey of UK general practice. *British Medical Journal* [online] 2(3) Available from: <http://bmjopen.bmj.com/content/2/3/e000851.full>

4] Gerald LB, Grad R, Turner-Henson A, Hains C, Tang S, Feinstein R, (2004) Validation of a multistage asthma case-detection procedure for elementary school children. *Pediatrics*.114(4) pp:459-68.

5] Jones CA, Morpew T, Clement LT, Kimia T, Dyer M, Li M, (2004) A school-based case identification process for identifying inner city children with asthma: the Breathmobile program. *Chest* 125(3) pp:924-34.

Cardiovascular data



West Somerset



| Significance | Indicator | West Somerset number | West Somerset value | Somerset average | Worst Locality value | Locality range <-Worse Better-> | Best Locality value | West Somerset value 2015 | Direction of change |
|--|---|----------------------|---------------------|------------------|----------------------|------------------------------------|---------------------|--------------------------|---------------------|
| | | | | | | | | | |
| Prevalence | 1 : Coronary heart disease | 1,745 | 39.0 | 37.6 | 42.8 | | 34.8 | 39.3 | |
| | 2 : Stroke/TIA | 1,378 | 30.9 | 27.3 | 30.9 | | 25.5 | 30.7 | |
| | 3 : Heart failure | 370 | 8.0 | 8.8 | 10.1 | | 8.0 | 7.6 | |
| | 4 : Atrial Fibrillation | 1,191 | 26.1 | 24.6 | 26.1 | | 23.8 | 25.5 | |
| | 5 : Hypertension | 6,479 | 150 | 161 | 176 | | 150 | 150 | |
| | 6 : Diabetes (ages 17 and over) | 2,091 | 61 | 66 | 75 | | 61 | 60 | |
| | 7 : Obesity (ages 16 and over) | 3,307 | 105 | 88 | 105 | | 73 | 88 | |
| Smoking | 8 : Current smokers (aged 16 and over) | | 13.2% | 12.8% | 17.8% | | 10.4% | 14.4% | |
| | 9 : 4 week smoking quit rate (all ages) | | 58% | 49% | 40% | | 60% | 43% | |
| | 10 : 4 week smoking quit rate (45-64) | | 42% | 45% | 30% | | 66% | 45% | |
| Mortality | 11 : Smokers going through cessation per 1000 recorded smokers | | 35 | 40 | 30 | | 54 | 81 | |
| | 12 : All circulatory disease | 686 | 103% | 100% | 110% | | 92% | 102% | |
| Admissions | 13 : All circulatory disease <75 | 117 | 97% | 100% | 124% | | 78% | 90% | |
| | 14 : Emergency admissions for Circulatory diseases (all ages) | 456 | 10.6 | 10.9 | 11.4 | | 9.4 | 9.0 | |
| | 15 : Elective admissions for Circulatory diseases (all ages) | 356 | 8.8 | 8.5 | 9.3 | | 6.9 | 8.4 | |
| | 16 : Emergency admissions for CHD | 254 | 2.0 | 2.3 | 2.6 | | 2.0 | 2.1 | |
| | 17 : Emergency admissions for CVD | 346 | 2.5 | 2.1 | 2.5 | | 1.8 | 2.5 | |
| NHS Health Checks | 18 : Alcohol related admissions | 3,650 | 1,833 | 1,894 | 2,062 | | 1,685 | 1,802 | |
| | 19 : Alcohol specific admissions | 849 | 495 | 445 | 548 | | 370 | 490 | |
| | 20 : % of eligible population checked | 243 | 11% | 17% | 8% | | 36% | 22% | |
| QOF 2015 ongoing management indicators | 21 : % of eligible population checked in most deprived quintile | 94 | 6% | 14% | 5% | | 44% | 14% | |
| | 22 : AF patients with CHA2DS2-VASc score measured | 891 | 97% | 84% | 71% | | 97% | | New indicators |
| | 23 : Anti-coagulation drug therapy in those AF patients with CHA2DS2-VASc score ≥ 2 | 789 | 82% | 84% | 80% | | 90% | | |
| | 24 : BP measured in patients ≥ 45 | 17,899 | 90% | 89% | 87% | | 92% | | |
| | 25 : CHD patients with 150/90 mmHg or less | 1,494 | 89% | 85% | 78% | | 95% | 90% | |
| | 26 : CHD patients with Aspirin or alternative taken | 1,566 | 93% | 91% | 88% | | 97% | | New indicator |
| | 27 : CHD patients had influenza immunisation | 1,310 | 94% | 89% | 84% | | 98% | 92% | |
| | 28 : HF patients currently treated with ACE-I or ARB | 39 | 95% | 87% | 80% | | 100% | 92% | |
| | 29 : HF patients currently treated with ACE-I or ARB also treated with beta-blocker | 30 | 94% | 86% | 80% | | 94% | 92% | |
| | 30 : Hypertension patients with 150/90 mmHg or less | 5,178 | 82% | 77% | 72% | | 86% | | New indicators |
| | 31 : PAD patients with 150/90 mmHg or less | 322 | 86% | 79% | 74% | | 92% | | |
| | 32 : PAD patients with Aspirin or alternative taken | 264 | 90% | 84% | 79% | | 94% | | |
| | 33 : Stroke/TIA patients with 150/90 mmHg or less | 922 | 87% | 80% | 76% | | 87% | 88% | |
| | 34 : NH Stroke/history of TIA patients taking antiplatelet agent or anti-coagulant | 608 | 95% | 91% | 88% | | 98% | 93% | |
| | 35 : Stroke/TIA patients had influenza immunisation | 810 | 94% | 88% | 81% | | 97% | 92% | |

| Indicator | Notes |
|-----------|---|
| 1-7 | MIQUEST (QOF) indirectly standardised prevalence rate within Somerset 2015 |
| 8-11 | Smoking prevalence from MIQUEST query March 2016 and SmokefreeLife Somerset service data April 2015 to March 2016 |
| 12-13 | Indirectly standardised mortality ratio (compared to Somerset) : ONS : 2011-15 |
| 14-19 | Indirectly standardised admission to hospital rate per 100,000 : Secondary Uses Service (SUS) : 2015/16 for Circulatory diseases, 2013/14 - 2015/16 for CHD and CVD, 2011/12 - 2015/16 for Alcohol admissions. Circulatory diseases ICD10 codes: Chapter I CHD ICD10 codes: I20-I25 CVD ICD10 codes: I6* |
| 20-21 | Alcohol related and alcohol specific causes as listed in work by North West Public Health Observatory and released through the Local Alcohol Profiles for England http://www.lape.org.uk/ |
| 22-35 | Somerset NHS Health Checks : financial year 2015/16 QOF ongoing management indicators : 2015 |

AF006: The percentage of patients with atrial fibrillation in whom stroke risk has been assessed using the CHA2DS2-VASc score risk stratification scoring system in the preceding 12 months (excluding those patients with a previous CHADS2 or CHA2DS2-VASc score of 2 or more)

AF007: In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anti-coagulation drug therapy

BP002: The percentage of patients aged 45 or over who have a record of blood pressure in the preceding 5 years

CHD002: The percentage of patients with coronary heart disease in whom the last blood pressure reading (measured in the preceding 12 months) is 150/90 mmHg or less

CHD005: The percentage of patients with coronary heart disease with a record in the preceding 12 months that aspirin, an alternative anti-platelet therapy, or an anti-coagulant is being taken

CHD007: The percentage of patients with coronary heart disease who have had influenza immunisation in the preceding 1 August to 31 March

HF003: In those patients with a current diagnosis of heart failure due to left ventricular systolic dysfunction, the percentage of patients who are currently treated with an ACE-I or ARB

HF004: In those patients with a current diagnosis of heart failure due to left ventricular systolic dysfunction who are currently treated with an ACE-I or ARB, the percentage of patients who are additionally currently treated with a beta-blocker licensed for Heart failure

HYP006: The percentage of patients with hypertension in whom the last blood pressure reading (measured in the preceding 12 months) is 150/90 mmHg or less

PAD002: The percentage of patients with peripheral arterial disease in whom the last blood pressure reading (measured in the preceding 12 months) is 150/90 mmHg or less

PAD004: The percentage of patients with peripheral arterial disease with a record in the preceding 12 months that aspirin or an alternative anti-platelet is being taken

STIA003: The percentage of patients with a history of stroke or TIA in whom the last blood pressure reading (measured in the preceding 12 months) is 150/90 mmHg or less

STIA007: The percentage of patients with a stroke shown to be non-haemorrhagic, or a history of TIA, who have a record in the preceding 12 months that an antiplatelet agent, or an anti-coagulant is being taken

STIA009: The percentage of patients with stroke or TIA who have had influenza immunisation in the preceding 1 August to 31 March

Diabetes profile



West Somerset

| | | | | | | | |
|--|--|--|---|--|---|--|-------------------------|
| | Significantly better than Somerset average | | Not significantly different to Somerset average | | Significantly worse than Somerset average | | Difference not assessed |
|--|--|--|---|--|---|--|-------------------------|

Worst Locality value

Chosen Locality

Best Locality

Other Localities

Somerset average

Improved

Worsened

| Significance | Indicator | West Somerset number | West Somerset value | Somerset average | Worst Locality value | Locality range ←Worse Better→ | Best Locality value | West Somerset value 2015 | Direction of change |
|--------------------------|--|----------------------|---------------------|------------------|----------------------|----------------------------------|---------------------|--------------------------|---------------------|
| Prevalence | 1 Diabetes (ages 17 and over) standardised rate | 2,091 | 61 | 66 | 75 | | 61 | 60 | |
| | 2 Diabetes (ages 17 and over) crude rate | 2,082 | 76 | 67 | 76 | | 58 | 70 | |
| | 3 % of diabetes cases on register | 2,047 | 62% | 65% | 73% | | 59% | 76% | |
| | 4 Obesity (ages 16 and over) | 3,307 | 104.6 | 87.8 | 104.6 | | 72.7 | 87.5 | |
| | 5 Excess weight (ages 4-5) | n/a | 22% | 22% | 26% | | 20% | 24% | |
| | 6 Excess weight (ages 10-11) | n/a | 30% | 30% | 32% | | 28% | 29% | |
| Deaths | 7 Diabetes as underlying cause | 32 | 0.14 | 0.11 | 0.18 | | 0.09 | 0.13 | |
| | 8 % of eligible checked | n/a | 11% | 17% | 8% | | 36% | 22% | |
| Health Checks | 9 % of eligible in most deprived quintile checked (capped at 100%) | n/a | 6% | 14% | 5% | | 44% | 14% | |
| | 10 Blood glucose lowering drugs | 15,624 | 83.5 | 79.7 | 77.4 | | 83.5 | 84.3 | |
| Prescribing | 11 Long/Intermediate Insulin Analogues | 827 | 53.6 | 67.7 | 83.4 | | 53.6 | 54.5 | |
| | 12 Emergency admissions for Diabetes (all ages) main dx | 178 | 0.76 | 0.95 | 1.21 | | 0.74 | 0.87 | |
| Admissions | 13 Emergency admissions (all ages) for someone with diabetes | 598 | 13.9 | 14.9 | 19.6 | | 10.8 | 13.0 | |
| | 14 Elective admissions (all ages) for someone with diabetes | 651 | 15.3 | 16.4 | 20.1 | | 12.6 | 14.8 | |
| | 15 Admissions (all ages) for lower limb amputation | 66 | 0.26 | 0.18 | 0.26 | | 0.14 | 0.23 | |
| | 16 Admissions (all ages) for DKA/coma | 63 | 0.32 | 0.26 | 0.55 | | 0.19 | 0.30 | |
| QOF 2015 care indicators | 17 DM002: BP<=150/90 mmHg | 1,796 | 91% | 86% | 80% | | 95% | 90% | |
| | 18 DM003: BP<=140/80 mmHg | 1,468 | 76% | 70% | 58% | | 83% | 75% | |
| | 19 DM004: Total cholesterol <=5 mmol/l | 1,460 | 81% | 79% | 76% | | 85% | 80% | |
| | 20 DM006: Patients with nephropathy/microalbuminuria given ACE | 165 | 90% | 84% | 78% | | 99% | 87% | |
| | 21 DM007: IFCC-HbA1c is <=59 mmol/mol | 1,301 | 71% | 64% | 55% | | 74% | 67% | |
| | 22 DM008: IFCC-HbA1c is <=64 mmol/mol | 1,466 | 78% | 73% | 66% | | 82% | 74% | |
| | 23 DM009: IFCC-HbA1c is <=75 mmol/mol | 1,704 | 89% | 84% | 79% | | 91% | 83% | |
| | 24 DM012: Had foot examination and risk classification | 1,771 | 90% | 81% | 73% | | 92% | 83% | |
| | 25 DM014: Been referred to a structured education programme | 93 | 86% | 74% | 54% | | 99% | 94% | |
| | 26 DM018: Had Flu vaccination | 1,485 | 94% | 89% | 84% | | 98% | 91% | |

- Indicator Notes
- 1, 4 MICEST (QOF) indirectly standardised prevalence rate per 1000 within Somerset 2016
 - 2 QOF crude prevalence rate per 1000, 2015
 - 3 Modelled true prevalence using York and Humber PHO models compared to QOF reported cases 2015
 - 5-6 National Child Measurement Programme 2010/11 - 2015/16; LSOA weighted average using population as at April 2016
 - 7 Indirectly standardised mortality ratio (compared to Somerset) : ONS : 2011-15
 - 8-9 Somerset NHS Health Checks : financial year 2015/16
 - 10-11 MO KTT prescribing indicators: NHS Business Authority report 2015/16
 - 12-16 Indirectly standardised admission to hospital (rate per 1000) : Secondary Uses Service (SUS) : 2015/16 for All emergency/elective admissions for those with diabetes; 2010/11 - 2015/16 for Diabetes as main reason for emergency admission, Lower limb amputations and DKA/coma admissions, Diabetes ICD10 codes: E10-E14; Lower limb amputations ICD10 codes E10-E14 AND OPCS4 codes X9-X11; DKA/Coma ICD10 codes E10.0, E10.1, E11.0, E11.1, E12.0, E12.1, E13.0, E13.1, E14.0, E14.1
 - 17-26 QOF ongoing management indicators : 2015

DM002: The percentage of patients with diabetes, on the register, in whom the last blood pressure reading (measured in the preceding 12 months) is 150/90 mmHg or less, NICE 2010 menu ID: NM01

DM003: The percentage of patients with diabetes, on the register, in whom the last blood pressure reading (measured in the preceding 12 months) is 140/80 mmHg or less, NICE 2010 menu ID: NM02

DM004: The percentage of patients with diabetes, on the register, whose last measured total cholesterol (measured within the preceding 12 months) is 5 mmol/l or less

DM006: The percentage of patients with diabetes, on the register, with a diagnosis of nephropathy (clinical proteinuria) or micro-albuminuria who are currently treated with an ACE-I (or ARBs)

DM007: The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 59 mmol/mol or less in the preceding 12 months, NICE 2010 menu ID: NM14

DM008: The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 64 mmol/mol or less in the preceding 12 months

DM009: The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 75 mmol/mol or less in the preceding 12 months

DM012: The percentage of patients with diabetes, on the register, with a record of a foot examination and risk classification: 1) low risk (normal sensation, palpable pulses), 2) increased risk (neuropathy or absent pulses), 3) high risk (neuropathy or absent pulses plus deformity or skin changes in previous ulcer) or 4) ulcerated foot within the preceding 12 months, NICE 2010 menu ID: NM13

DM010: The percentage of patients with diabetes, on the register, who have had influenza immunisation in the preceding 1 September to 31 March

DM014: The percentage of patients newly diagnosed with diabetes, on the register, in the preceding 1 April to 31 March who have a record of being referred to a structured education programme within 9 months after entry on to the diabetes register, NICE 2011 menu ID: NM27

DM018: The percentage of patients with diabetes, on the register, who have had influenza immunisation in the preceding 1 August to 31 March (the comparative data for previous year is the same but from September to March)

Cancer Profile

West Somerset



| Indicator | West Somerset number | West Somerset value | Somerset CCG value | Lowest Locality value in Somerset | Chart | Highest Locality value in Somerset | West Somerset compared to Somerset | England value | West Somerset previous time period | Direction of change |
|--|----------------------|---------------------|--------------------|-----------------------------------|-------|------------------------------------|------------------------------------|---------------|------------------------------------|---------------------|
| 1 Cancer: QOF prevalence (all ages) (%) | 1,393 | 4.4 | 3.2 | 2.7 | | 4.4 | Higher than Somerset | 2.4 | 4.0 | Increased |
| 2 New cancer cases (Crude incidence rate: new cases per 100,000 population) | 258 | 821 | 602 | 511 | | 821 | Higher than Somerset | 515 | 823 | Decreased |
| 3 % reporting cancer in the last 5 years | 19 | 4.1 | 3.7 | 2.9 | | 4.6 | Higher than Somerset | 3.2 | 5.9 | Decreased |
| 4 CAN003: review within 6 mths of diagnosis (%) | 152 | 71 | 46 | 31 | | 71 | Higher than Somerset | 71 | 83 | Decreased |
| 5 CS002: Women, aged 25-64, with a record of cervical screening (last 5 yrs) (%) | 5,159 | 71 | 76 | 71 | | 79 | Higher than Somerset | 76 | 73 | Decreased |
| 6 Females, 25-64, attending cervical screening within target period (3.5 or 5.5 year coverage, %) | 4,849 | 71 | 75 | 71 | | 78 | Higher than Somerset | 74 | 72 | Decreased |
| 7 Females, 50-70, screened for breast cancer in last 36 months (3 year coverage, %) | 3,887 | 73 | 75 | 69 | | 78 | Higher than Somerset | 73 | 75 | Decreased |
| 8 Females, 50-70, screened for breast cancer within 6 months of invitation (Uptake, %) | 127 | 49 | 76 | 41 | | 80 | Higher than Somerset | 74 | 47 | Increased |
| 9 Persons, 60-69, screened for bowel cancer within 6 months of invitation (Uptake, %) | 1,656 | 63 | 63 | 61 | | 65 | Higher than Somerset | 58 | 57 | Increased |
| 10 Persons, 60-69, screened for bowel cancer in last 30 months (2.5 year coverage, %) | 3,328 | 62 | 62 | 60 | | 64 | Higher than Somerset | 58 | 61 | Increased |
| 11 In-patient or day-case colonoscopy procedures (Number per 100,000 population) | 203 | 638 | 640 | 572 | | 743 | Higher than Somerset | 733 | 546 | Increased |
| 12 In-patient or day-case sigmoidoscopy procedures (Number per 100,000 population) | 317 | 997 | 816 | 418 | | 1,025 | Higher than Somerset | 478 | 830 | Increased |
| 13 In-patient or day-case upper GI endoscopy procedures (Number per 100,000 population) | 453 | 1,424 | 1,338 | 1,164 | | 1,539 | Higher than Somerset | 1,302 | 1,315 | Increased |
| 14 Number of emergency admissions with cancer (Number per 100,000 population) | 303 | 953 | 667 | 575 | | 953 | Higher than Somerset | 538 | 782 | Increased |
| 15 Number of emergency presentations (Number per 100,000 population) | 59 | 186 | 116 | 96 | | 186 | Higher than Somerset | 89 | 123 | Increased |
| 16 Number of other presentations (Number per 100,000 population) | 178 | 560 | 422 | 321 | | 560 | Higher than Somerset | 361 | 464 | Increased |
| 17 Two-week wait referrals for suspected cancer (Number per 100,000 population) | 1,212 | 3,811 | 3,394 | 2,925 | | 3,842 | Higher than Somerset | 2,975 | 3,318 | Increased |
| 18 Two-week wait referrals (Indirectly age-sex standardised referral ratio) | 1,212 | 93 | 99 | 92 | | 105 | Higher than Somerset | 100 | 89 | Increased |
| 19 Two-week referrals resulting in a diagnosis of cancer (Conversion rate: as % of all TWW referrals). | 151 | 12.5 | 10.0 | 7.7 | | 12.5 | Higher than Somerset | 7.8 | 9.2 | Increased |
| 20 Number of new cancer cases treated (Detection rate: % of which resulted from a TWW referral) | 152 | 56 | 53 | 47 | | 64 | Higher than Somerset | 50 | 45 | Increased |
| 21 Two-week wait referrals for suspected breast cancer (Number per 100,000 population) | 216 | 679 | 535 | 396 | | 723 | Higher than Somerset | 541 | 549 | Increased |
| 22 Two-week wait referrals for suspected lower GI cancers (Number per 100,000 population) | 195 | 613 | 502 | 426 | | 613 | Higher than Somerset | 453 | 561 | Increased |
| 23 Two-week wait referrals for suspected lung cancer (Number per 100,000 population) | 49 | 154 | 107 | 81 | | 154 | Higher than Somerset | 103 | 129 | Increased |
| 24 Two-week wait referrals for suspected skin cancer (Number per 100,000 population) | 220 | 692 | 735 | 572 | | 869 | Higher than Somerset | 572 | 536 | Increased |
| 25 Two-week wait referrals for suspected cancer (Number per 100,000 population). Five years combined data. | 5,171 | 16,259 | 14,381 | 11,992 | | 16,450 | Higher than Somerset | 11,860 | 15,237 | Increased |
| 26 Two-week wait referrals (Indirectly age-sex standardised referral ratio). Five years combined data. | 5,171 | 98 | 105 | 91 | | 113 | Higher than Somerset | 100 | 101 | Decreased |
| 27 Two-week referrals resulting in a diagnosis of cancer (Conversion rate: as % of all TWW referrals). Five years combined data. | 628 | 12.1 | 10.8 | 8.1 | | 12.1 | Higher than Somerset | 8.8 | 12.6 | Decreased |
| 28 Number of new cancer cases treated (Detection rate: % of which resulted from a TWW referral). Five years combined data. | 649 | 51 | 52 | 43 | | 54 | Higher than Somerset | 48 | 48 | Increased |
| 29 Two-week wait referrals for suspected breast cancer (Number per 100,000 population). Five years combined data. | 822 | 2,585 | 2,252 | 1,636 | | 2,966 | Higher than Somerset | 2,161 | 2,325 | Increased |
| 30 Two-week wait referrals for suspected lower GI cancers (Number per 100,000 population). Five years combined data. | 936 | 2,943 | 2,280 | 1,987 | | 2,943 | Higher than Somerset | 1,897 | 2,782 | Increased |
| 31 Two-week wait referrals for suspected lung cancer (Number per 100,000 population). Five years combined data. | 191 | 601 | 477 | 390 | | 605 | Higher than Somerset | 449 | 577 | Increased |
| 32 Two-week wait referrals for suspected skin cancer (Number per 100,000 population). Five years combined data. | 774 | 2,434 | 2,828 | 2,395 | | 3,358 | Higher than Somerset | 2,170 | 2,094 | Increased |

Notes

- (1) Current data is for time periods ending in 2015/16 and previous data is for time periods ending in 2014/15
- (2) Data from PHE National General Practice Profiles <https://fingertips.phe.org.uk/profile/general-practice/data#mod.2.pyr.2016.pat.19.par.E38000150.are.L85005.sid1.2000005.ind1-.ind2.->

The following information is presented:

- Deaths – overview of causes of death 2011-2015 of Commissioning Locality patients.
- Years of Life Lost – overview of causes of years of life lost due to death before age 75 2011-2015 in Commissioning Locality patients.
- Life expectancy
- Standardised Mortality Ratios
- Where people die

The 'blobs' diagrams were first introduced in the Somerset 2013 profiles and provide an at a glance indication of the main causes of death 2011-2015 in Commissioning Locality registered patients. Different colours represent different disease/condition groups and lines indicate how more specific conditions are included within a larger grouping. The two 'blobs' with no lines have overlapping content with other 'blobs' but are not strict subgroups. They are placed beside the mental and behavioural disorders as they are, at least in part, mental health issues.

Possibly more important in terms of public health impact and valuable interventions is the corresponding diagram showing the conditions leading to years of life lost due to death before the age of 75. Whilst circulatory diseases cause approximately a third of deaths, they account for less than 20% of the years of life lost. With regard to years of life lost (YLL), cancer is the main cause and justifiably receives a lot of attention, with lung cancer being of particular concern. Other cancers which result in a high proportion of years lost have corresponding screening programmes – breast cancer, colorectal cancer and if YLL are high it is important to ensure that screening programmes are working as effectively as possible.


Life expectancy is presented at birth and also additional life expectancy at age 65. On the whole Somerset life expectancy is longer than England as a whole. Another index of preventable deaths is the proportion of deaths occurring before the age of 65 and Somerset is also shown to be better than the national average.

Standardised mortality ratios look at the number of deaths compared to the number expected if the Commissioning Locality followed the average Somerset age/sex specific death rates. They can highlight specific causes of death which may be more prevalent in this Commissioning Locality. Of particular concern are the causes of death occurring at young ages and robbing people of many valuable years of life. 75 years is often used as an arbitrary age to highlight the issue of premature deaths, without implying that years of life lived after that age are in any way less valuable.

Where people die can give clues as to care available. Research has shown that many people with terminal conditions prefer to die at home if they have the choice and are appropriately supported. Numbers of those dying in nursing homes in part reflects location of nursing homes and any local barriers or enablers to access. Lower rates of people dying in hospices could also reflect access issues. High numbers of those dying in hospital could be due to emergency admissions for disease that has not been well managed. More detail on hospitalisations is given in the later section on admissions.


It is worth noting that ONS have revised mortality coding twice in the time period covered by these statistics - in 2011 and in 2014 - to be in line with how cause of death is coded internationally. This has had some impact on the proportion of deaths coded to certain conditions, notably dementia and diabetes. Further details of the effects can be found at http://www.ons.gov.uk/ons/dcp171778_373602.pdf.

DEATHS



*There is some overlap between Drug related deaths and Mental and behavioural disorders and also between Drug related deaths and Suicide and injury undetermined deaths. There is a further overlap between Drug related deaths and Accidents.

YEARS OF LIFE LOST BEFORE THE AGE OF 75




*There is some overlap between Drug related deaths and Mental and behavioural disorders and also between Drug related deaths and Suicide and injury undetermined deaths. There is a further overlap between Drug related deaths and Accidents.

Life expectancy 2011-15


Life expectancy based on deaths from ONS Primary Care Mortality Database and population estimates from the Exeter system

| | Commissioning Locality | Somerset | England (2012-14) | Range of Practice values low / median / high |
|--|------------------------|----------|-------------------|--|
| Male life expectancy at birth | 81.6 | 81.1 | 79.5 | 76.0 / 81.5 / 86.1 |
| Female life expectancy at birth | 84.9 | 84.5 | 83.2 | 80.8 / 85.1 / 96.1 |
| Male life expectancy at 65 | 20.4 | 19.6 | 19.5 | 15.4 / 19.9 / 31.0 |
| Female life expectancy at 65 | 22.8 | 22.0 | 22.0 | 18.8 / 22.4 / 33.8 |
| Proportion of deaths occurring before age 65 | 10.7% | 12.0% | 15.9% | 5.9% / 12.0% / 66.0% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of life expectancy of the resident population. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show shorter life expectancy compared to other Commissioning Localities, values to the right show longer life expectancy and so better performance. For the proportion of deaths occurring before 65 a score to the left indicates more deaths prior to age 65 compared to others.



The Commissioning Locality has the best value in the county for:
Female life expectancy at 65




Standardised mortality ratios

Deaths registered between 2011 and 2015, age and sex standardised to Somerset as a whole (a value of 100% is the Somerset average). ONS Primary Care Mortality Database.

The standardised mortality ratio (SMR) describes the mortality rate in the adjusting for differences between Commissioning Localities in their age and sex profile and compared to the overall mortality in Somerset. A ratio greater than 100% indicates higher rates of death than expected, a ratio of 110% indicates death rates 10% higher than in Somerset as a whole. Where the all causes, all ages SMR is high, the other SMRs presented for more specific ages and/ or causes may indicate the root of the imbalance or specific issues which may be masked in the overall SMR.

| Condition | Observed | Expected | Commissioning Locality SMR | Somerset SMR | England | Range of Practice values low / median / high |
|------------------------------------|----------|----------|----------------------------|--------------|---------|--|
| All causes - all ages | 2,150 | 2,312 | 93% | 100% | | 71% / 97% / 142% |
| All causes - <75 | 542 | 542 | 100% | 100% | | 59% / 97% / 187% |
| All causes - 85+ | 1,022 | 1,094 | 93% | 100% | | 29% / 96% / 156% |
| Cancer - all ages | 637 | 624 | 102% | 100% | | 73% / 101% / 156% |
| Cancer - <75 | 276 | 249 | 111% | 100% | | 67% / 102% / 208% |
| All circulatory disease - all ages | 686 | 668 | 103% | 100% | | 56% / 95% / 147% |
| All circulatory disease - <75 | 117 | 120 | 97% | 100% | | 39% / 95% / 209% |
| Coronary heart disease - all ages | 319 | 280 | 114% | 100% | | 51% / 95% / 194% |
| All respiratory disease - all ages | 223 | 289 | 77% | 100% | | 53% / 96% / 141% |
| COPD - all ages | 81 | 115 | 71% | 100% | | 42% / 101% / 305% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of SMRs. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. For consistency with previous graphs, values to the left show a worse position, which equates to a greater SMR. Values to the right show lower SMRs compared to other Commissioning Localities.



The Commissioning Locality has the worst value in the county for:





Cancer - <75 Coronary heart disease - all ages

Significantly better than county average for:

All causes - all ages All respiratory disease - all ages COPD - all ages

The Commissioning Locality has the best value in the county for:

All respiratory disease - all ages COPD - all ages



West Somerset


Where people die

Deaths registered between 2011 and 2015. ONS Primary Care Mortality Database.


Where people die will reflect access to services as well as patterns of care. The category of Anywhere else covers everywhere not in the other categories and includes such places as other people's houses and roads.

| | Commissioning Locality % | Somerset % | England % | Range of Practice values low / median / high |
|-------------------|--------------------------|------------|-----------|--|
| Home | 24% | 22% | 22% | 9% / 24% / 40% |
| Care/nursing home | 27% | 28% | 21% | 5% / 26% / 53% |
| Hospital | 43% | 42% | 48% | 27% / 43% / 56% |
| Anywhere else | 6% | 8% | 8% | 3% / 8% / 18% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. For the purposes of this data, it is assumed that it is better to die at home and worse to die anywhere else. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show a worse position and values to the right show a better position compared to other Commissioning Localities.



* for this diagram it is assumed that it is better to die at home and worse to die anywhere else.



The profile contains information on the following preventative health aspects which may impact on the local Commissioning Locality.

The following information is presented:

- Screening, health checks, cancer screening, chlamydia
- Immunisations, childhood and seasonal influenza
- Breastfeeding – initiation rates and continuation at 6-8 weeks

The Somerset NHS Health Check programme was recommissioned by Somerset County Council in April 2015. The health checks are delivered through GP practices, pharmacies, workplaces, community venues and through the use of a mobile van. More detail on this programme is provided in the health checks profiles for each Commissioning Locality which highlights checks taken up and the coverage by socioeconomic indicators as well as outcomes.

The outcomes of three cancer screening programmes are shown: bowel cancer, breast cancer, and cervical screening. Data is shown for the current year and previous year. For the cervical screening programme, the rate of inadequate smears is also provided.

Chlamydia screening is provided at a number of locations within the county so overall rates of screening are higher than just those screened via the GP. However local research suggests that young people are very receptive to screening provided routinely in general practice so the overall low screening rates probably represent a missed opportunity to improve health. More detail on interventions which can support increased chlamydia screening is provided on page 36.

The goal is for coverage for all antigens in the childhood vaccination schedule to be at 95% as this is a level which guarantees herd immunity. A selection of childhood vaccinations is presented. In general Somerset childhood immunisation rates are similar to or better than those seen in the rest of England.

A new Children and Young People's Health Services (CYPHS) dataset was introduced in April 2015 but only contains data from September 2015. Currently there is not a full year's robust data for breastfeeding, so data has been presented for 2014/15 in this profile. The initiation rates in Somerset are, in general, better than those seen in the rest of England although there can be quite striking variation by practice. Whilst GPs may not be so close to the routine support provided to mothers on breastfeeding, they may be consulted about problems which may lead to earlier cessation of breastfeeding than mothers may ideally choose.

The paediatric profile provides an overview of many issues which may impact on the health of children, including breastfeeding, immunisation, childhood obesity, teenage deliveries, youth admission rates and wider determinants of health. This profile was developed to support a Somerset CCG study day raising the profile of child health needs and highlighting the role of GPs in this - for example promoting healthy weight and being aware of the health impact of children living in households with a low income. The aim is to provide a more holistic approach to childhood health.

NHS Health Checks - notes to accompany activity snapshot

Local Authorities have a statutory duty to invite their eligible population for a NHS Health Check once every 5 years as part of the 5-year call/recall. In addition, local authorities have a legal duty to seek continuous improvement in the percentage of eligible individuals taking up their offer of a NHS Health Check as part of their statutory duties. As of January 2016, there were 261,000 people in Somerset aged 40 to 74 of whom 185,325 (71%) were eligible for a NHS Health Check every 5 years.

The Somerset NHS Health Check programme was recommissioned by Somerset County Council in April 2015. The health checks are delivered through GP practices, pharmacies, workplaces, community venues and through the use of a mobile van.

CHART OF ELIGIBLE POPULATION & HEALTH CHECKS COMPLETED

The charts indicate levels of population activity by Commissioning Locality and by County level:

- 1) The proportion of the total population aged 40-74 registered with a GP Practice during the current year (20%).
- 2) The proportion of individuals aged 40 to 74 (from item 1 above) who are eligible for a health check as they have not been diagnosed by their GP with a pre-existing condition.
- 3) The proportion of eligible individuals (from item 2 above) who have received a NHS Health Check

CHART OF COMPLETED CHECKS BY DEPRIVATION QUINTILES

These charts depict health check activity within each deprivation quintile.

- 1) Indicates the number of people in each quintile eligible for a check and the number of checks completed in that quintile, for the Commissioning Locality.
- 2) Indicates the checks completed as a percentage of the total eligible population in each quintile of deprivation, for the Commissioning Locality.
- 3) Indicates the checks completed as a percentage of the total eligible population in each quintile of deprivation, for the Commissioning Locality.

TABLE OF NHS HEALTH CHECKS ACTIVITY

The NHS Health Check data is categorised by Activity, Health Check Results and Referrals.

Columns marked Locality Lowest & Locality Highest: Values of 0% or 100% may appear within these 2 categories where only a few checks have been completed for the Commissioning Locality.

TARGET ACTIVITY: Indicates the number and percentage of people eligible for a health check and who received a health check.

HEALTH CHECK RESULTS: This section indicates the findings from completed health checks. The following criteria are new to the Somerset NHS Health Check programme:


- Item 13 Diabetes Filter triggered: Where a person's BP is recorded as $\geq 140/90$ and/or their BMI is ≥ 30 , this indicates the possibility of a risk of diabetes.
- Item 14 raised HbA1c: HbA1c test was available for health checks delivered in workplaces or community locations.
- Item 15 Family History of CVD: This is recorded when the person indicates this is the case for a family member in the 1st degree.
- Item 16 Depression Assessment.

LIFESTYLE COUNSELLING AND REFERRALS: This section indicates where there was a recommendation that the user should follow up with their GP or where the user requested referral to a lifestyle support service.


Referral activity percentages are calculated against total checks completed EXCEPT where the referral is based on a specific health check result (eg number of smokers identified who requested a referral to the stop smoking service).

West Somerset


The population/activity charts below provide a visual representation of NHS Health Check activity compared to the annual eligible population. This is by Commissioning Locality as at 2016 and Somerset County.



Numbers eligible and checked by deprivation



Checks as % of eligible by deprivation




West Somerset

West Somerset

SOMERSET

| Indicator | | Locality Number | Locality Value | Somerset average | Lowest Locality | Range of Locality values | Highest Locality | |
|-------------------------------------|--|---|----------------|------------------|-----------------|--------------------------|------------------|-------|
| Target activity | 1: % eligible for check | 2,269 | 69% | 71% | 68% | | 74% | |
| | 2: Checks as % of eligible | 243 | 11% | 17% | 8% | | 36% | |
| | Health Check Results | 3: Current Smoker | 18 | 7% | 11% | 7% | | 16% |
| | | 4: Very High Alcohol score | 1 | 0.4% | 0.3% | 0.0% | | 0.5% |
| | | 5: Chest/Calf Pain indicated* | 3 | 1.2% | 2.2% | 1.2% | | 5.1% |
| | | 6: AF screening / Irregular pulse* | 5 | 2.1% | 1.8% | 0.4% | | 3.0% |
| | | 7: BP Raised ($\geq 140/90$ and $< 160/100$)* | 65 | 27% | 23% | 19% | | 30% |
| | | 8: BP High ($\geq 160/100$)* | 25 | 10.3% | 7.4% | 5.2% | | 10.3% |
| | | 9: GPPAQ = Moderately Inactive | 38 | 16% | 17% | 13% | | 21% |
| | | 10: GPPAQ = Inactive | 91 | 37% | 28% | 25% | | 37% |
| | | 11: Obese or morbidly obese (BMI ≥ 30) | 41 | 17% | 20% | 17% | | 26% |
| | | 12: Non-fasting Chol/HDL Ratio $\geq 7^*$ | 11 | 4.5% | 3.8% | 2.8% | | 4.8% |
| | 13: Diabetes Filter triggered* | 96 | 40% | 39% | 32% | | 44% | |
| | 14: HbA1c (Outreach**) Raised (≥ 42 mmol/mol)* | 5 | 8% | 12% | 8% | | 17% | |
| | 15: Family History of CVD | 95 | 39% | 38% | 37% | | 42% | |
| | 16: Depression (PRIME-MD) | 31 | 13% | 12% | 9% | | 17% | |
| | 17: Qrisk ≥ 20 (high & very high risk)* | 30 | 12.3% | 6.3% | 4.5% | | 12% | |
| Lifestyle Counselling and Referrals | 18: All GP Referrals | 126 | 52% | 48% | 41% | | 54% | |
| | 19: Smokers referred to smoking cessation | 1 | 6% | 14% | 5% | | 21% | |
| | 20: Referrals for Very High Alcohol score | 0 | 0% | 29% | 0% | | 100% | |
| | 21: Referral for mental wellbeing | 14 | 45% | 39% | 28% | | 45% | |
| | 22: Qrisk ≥ 20 (high & very high risk) | 30 | 12.3% | 6.3% | 4.5% | | 12% | |
| | 23: Total Referred PA GPPAQ = Moderately Inactive | 3 | 7.9% | 6.7% | 1.3% | | 14% | |
| | 24: Total Referred PA GPPAQ = Inactive | 7 | 7.7% | 8.3% | 2.6% | | 16% | |
| | 25: Referral for Weight Management | 9 | 22% | 32% | 20% | | 47% | |



* Requires referral to GP (NICE Guidance)
 ** Outreach includes Community and Workplace locations
 All people receive lifestyle counselling

Feedback should be directed to Sharon Ashton at seashton@somerset.gov.uk

Cancer screening

Breast cancer screening from KC63 report and Cervical cancer screening from KC53 report . Bowel cancer screening from NHS England. All provided by the CSU.

Of note is the often wide variation in screening rates at practice level and to aid further work within the Locality we have highlighted where the practices rank.


Bowel and Cervical 2015/16, Breast 2014/15

| Condition | Commissioning Locality | Somerset | England (2015 - ages 60-74 - for bowel) | Range of Practice values low / median / high |
|---|------------------------|----------|---|--|
| Bowel: 60-69 uptake of invitation to screen in year | 57.9% | 58.2% | 57.1% | 41% / 58% / 66% |
| Breast: ages 45-49: coverage | 13.7% | 12.6% | 66.7% | 0% / 3% / 43% |
| Breast: ages 50-52: coverage | 61.1% | 57.5% | 69.0% | 14% / 64% / 89% |
| Breast: ages 53-64: coverage | 76.4% | 77.3% | 71.3% | 39% / 79% / 85% |
| Breast: ages 65-70: coverage | 79.2% | 77.8% | 72.9% | 15% / 80% / 88% |
| Breast: ages 71-74: coverage | 55.2% | 51.9% | 68.4% | 7% / 56% / 82% |
| Cervical: ages 25-64 coverage within 5 years | 72.1% | 77.6% | 76.5% | 66% / 78% / 88% |
| Cervical: ages 25-49 coverage within 3.5 years | 66.5% | 72.3% | 70.3% | 60% / 73% / 81% |
| Cervical: ages 50-64 coverage within 5 years | 70.9% | 76.0% | 75.7% | 60% / 76% / 88% |
| Cervical: Inadequate smears (ages 20-64) | 1.6% | 2.1% | 2.5% | 0% / 2% / 7% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their coverage for cancer screening. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond.

For screening uptake indicators, values to the left show lower levels of screening and worse performance and values to the right show higher rates of screening and better performance.


For the inadequate smears indicator, values to the left indicate higher than average rates of inadequate smear tests and thus worse performance. Values to the right indicate lower relative rates of inadequate smears and better performance.



The Commissioning Locality has the worst value in the county for:
 Cervical: ages 25-64 coverage within 5 years Cervical: ages 25-49 coverage within 3.5 years Cervical: ages 50-64 coverage within 5 years

Previous year





| Condition | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|---|------------------------|----------|---------|--|
| Bowel: 60-69 uptake of invitation to screen in year | 61.3% | 62.0% | 57.1% | 40% / 62% / 68% |
| Breast: ages 45-49: coverage | 21.2% | 11.2% | 68.3% | 0% / 2% / 42% |
| Breast: ages 50-52: coverage | 75.6% | 54.6% | 70.5% | 17% / 66% / 82% |
| Breast: ages 53-64: coverage | 77.4% | 79.1% | 72.1% | 54% / 80% / 85% |
| Breast: ages 65-70: coverage | 79.1% | 80.3% | 73.4% | 59% / 80% / 88% |
| Breast: ages 71-74: coverage | 41.6% | 55.2% | 68.4% | 16% / 59% / 81% |
| Cervical: ages 25-64 coverage within 5 years | 73.8% | 78.4% | 77.2% | 68% / 79% / 87% |
| Cervical: ages 25-49 coverage within 3.5 years | 67.9% | 73.1% | 71.3% | 61% / 74% / 84% |
| Cervical: ages 50-64 coverage within 5 years | 72.9% | 77.4% | 76.5% | 63% / 78% / 88% |
| Cervical: Inadequate smears (ages 20-64) | 4.8% | 2.9% | 2.5% | 0% / 3% / 7% |



The Commissioning Locality has the best value in the county for:
 Breast: ages 45-49: coverage Breast: ages 50-52: coverage

The Commissioning Locality has the worst value in the county for:
 Breast: ages 71-74: coverage Cervical: ages 25-64 coverage within 5 years Cervical: ages 25-49 coverage within 3.5 years Cervical: ages 50-64 coverage within 5 years Cervical: Inadequate smears (ages 20-64)

Cancer screening - latest year



Chlamydia screening




Uptake of Chlamydia screening 2015/16

Chlamydia screening is provided at a number of locations around the county. The data shown is the proportion of eligible population aged 15-24 being screened for Chlamydia in practices.

2015/16

| | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|---|-----------------------------|---------------|---------|--|
| Uptake of Chlamydia screening in those aged 15-24 | 2.1% | 3.0% | | 0% / 2% / 26% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their coverage for screening over the past three years. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show lower levels of screening and worse performance and values to the right show higher rates of screening and better performance.





General practice is essential for ensuring young people aged 15-24 years have access to chlamydia screening, particularly in a rural county like Somerset. Consultations with young people in Somerset show that the many would like to be able to access sexual health services via general practice; a local survey showed that 50% would prefer to go to their surgery for a chlamydia test and that general practice was the most popular place for them to accept testing when offered

Somerset-wide Integrated Sexual Health (SWISH) services alongside the Somerset Chlamydia Screening Programme strive to increase the detection rate of Chlamydia in the region. The routine availability of chlamydia screening in general practice is an essential component of the strategy to reduce sexually transmitted infections in young people who carry one of the highest burdens of sexual ill health in Somerset.

Below are some of the tried and tested interventions in general practice known to improve the offer and uptake of chlamydia screening by young people:

- Make it normal, advise it is something that is routinely offered to young people as part of keeping them healthy and link with other health promotion messages
- Ensure that young people are clear that the test is easy to do, self-taken, not invasive and confidential
- Introduce opportunities not requiring direct nurse/doctor involvement – ensure posters and leaflets are displayed in waiting rooms and treatment rooms and leave kits in grab bins in reception, resources or toilets
- Advise that all young people should have a chlamydia test once a year and with every new partner
- Identify a member of staff as your 'chlamydia screening champion'
- The champion can ensure your surgery maximises every opportunity to promote chlamydia screening. Previous evidence from Somerset shows that those surgeries that nominated a champion saw a significant increase in their test returns e.g. one surgery has increased their screens from 37 a year to consistently achieving well over a hundred screens per year for the past 3 years
- Have a 'whole team' approach and make greater use of all staff in the surgery. This could include reception staff giving out information leaflets or test kits before the young person sees their clinician or the use of health care assistants who can see young people before or after their appointment
- Every time a young person visits their general practice is an opportunity to offer a chlamydia screen
- Tack the offer of a test onto specific clinics e.g. all sexual health and contraceptive clinics, new patient health checks and travel clinics
- Use a 'pop-up' on EMIS for everyone aged 15-24. It is useful to do this for all contraceptive appointments but including everyone ensures that young men are offered the test
- Put the emphasis on 'on site' completion of the test and immediate return – less chance of 'losing it', more chance of 'getting round' to it
- If a young person is with friends offer them all a test
- Make kits available to the partners of young people particularly if they have tested positive through their GP service

Please contact the Somerset Chlamydia Screening Office on 01749 836704 or Somersetcs@sompar.nhs.uk for


- more information on the Somerset Chlamydia Screening programme
- ordering resources such as kits, grab bins, posters, leaflets and window stickers
- arranging a visit by one of the team to advise on how to improve your offer of chlamydia screening

Seasonal flu vaccinations

Flu vaccination programme statistics. Winter 2015/16



| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|----------------------------|------------------------|----------|---------|--|
| Those aged 65 and over | 65% | 70% | 71% | 59% / 70% / 82% |
| All children aged 2,3 or 4 | 41% | 42% | 34% | 9% / 44% / 74% |
| Those aged <65 at risk | 42% | 46% | 45% | 34% / 46% / 64% |
| Pregnant women | 38% | 43% | 42% | 20% / 42% / 68% |
| Carers | 26% | 37% | n/a | 13% / 39% / 70% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their immunisation outcomes. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show lower relative levels of immunisation and so worse performance and values to the right show higher rates of immunisation and so better performance.



The Commissioning Locality has the worst value in the county for:

Those aged 65 and over Those aged <65 at risk





Childhood Immunisations

Data from NHS England : Child Immunisation Collection 2015/16 outturn - this is management information and may not be exactly the same as COVER

2015/16


| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|---|------------------------|----------|---------|--|
| 5-in-1 primary vaccination uptake by 1st birthday | 97.4% | 95.3% | 93.6% | 81% / 96% / 100% |
| 4-in-1 booster vaccination uptake by 5th birthday | 95.1% | 94.1% | 86.3% | 75% / 95% / 100% |
| MMR primary vaccination uptake by 2nd birthday | 95.0% | 95.0% | 91.9% | 73% / 96% / 100% |
| MMR2 second dose uptake by 5th birthday | 95.1% | 92.7% | 88.2% | 67% / 94% / 100% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their immunisation outcomes. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show lower relative levels of immunisation and so worse performance and values to the right show higher rates of immunisation and so better performance.







previous year

| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|---|------------------------|----------|---------|--|
| 5-in-1 primary vaccination uptake by 1st birthday | 96.0% | 95.5% | 94.2% | 78% / 96% / 100% |
| 4-in-1 booster vaccination uptake by 5th birthday | 93.2% | 93.8% | 88.5% | 74% / 95% / 100% |
| MMR primary vaccination uptake by 2nd birthday | 96.0% | 94.9% | 92.3% | 73% / 95% / 100% |
| MMR2 second dose uptake by 5th birthday | 92.8% | 92.9% | 88.6% | 74% / 93% / 100% |



Childhood immunisations - latest year



West Somerset

Breastfeeding initiation and prevalence at 6-8 weeks


During transition to new maternity dataset the Breastfeeding data available is not currently robust. 2014/15 data only is shown below. Breastfeeding initiation and prevalence at 6 to 8 weeks data from NHS England Statistical Release: 12 week Maternal Assessment, Breastfeeding Initiation & Breastfeeding prevalence 6-8 weeks at Practice level OT 2014/15

2014/15


| Condition | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|--|------------------------|----------|---------|--|
| Number of maternities | 233 | 5,431 | 634,378 | |
| Percentage mothers initiating breastfeeding* | 88% | 81% | 74% | 60% / 82% / 100% |
| Number of infants due a check at 6-8 weeks | 220 | 5,281 | 631,353 | |
| Percentage of infants being wholly or partially breastfed* | 52% | 49% | 44% | 30% / 50% / 92% |

* of those with known status

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities in terms of their breastfeeding outcomes. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show lower levels of breastfeeding and worse performance and values to the right show higher rates of breastfeeding and better performance.



The Commissioning Locality has the best value in the county for:
Percentage mothers initiating breastfeeding*



Paediatric Profile



West Somerset



| Indicator | West Somerset number | West Somerset value | Somerset CCG value | Worst Locality value in Somerset | Chart | Best Locality value in Somerset | England value | Worst practice value in Somerset | Best practice value in Somerset | West Somerset value 2015 | Direction of change |
|--|----------------------|---------------------|--------------------|----------------------------------|-------|---------------------------------|---------------|----------------------------------|---------------------------------|--------------------------|---------------------|
| 1 % of population aged 0-14 * | 4,005 | 12% | 16% | 18% | | 12% | 18% | 24% | 9% | 12% | Higher |
| 2 Income deprivation affecting Children Index | | 0.17 | 0.14 | 0.20 | | 0.13 | 0.15 | 0.26 | 0.06 | 0.17 | Lower |
| 3 Breastfeeding initiation | 204 | 88% | 81% | 75% | | 88% | 74% | 60% | 100% | 88% | Improved |
| 4 Breastfeeding prevalence at 6-8 weeks | 115 | 52% | 49% | 40% | | 54% | | 30% | 92% | 52% | Improved |
| 5 5-in-1 uptake by age 1 | 226 | 97% | 95% | 91% | | 98% | 94% | 81% | 100% | 96% | Improved |
| 6 4-in-1 booster uptake by age 5 | 251 | 95% | 94% | 91% | | 97% | 86% | 75% | 100% | 93% | Improved |
| 7 MMR uptake by age 2 | 267 | 95% | 95% | 92% | | 96% | 92% | 73% | 100% | 96% | Improved |
| 8 MMR booster uptake by age 5 | 251 | 95% | 93% | 89% | | 96% | 88% | 67% | 100% | 93% | Improved |
| 9 Flu vaccination uptake for 2, 3 and 4yr olds | 340 | 41% | 42% | 32% | | 47% | 34% | 9% | 74% | 42% | Improved |
| 10 % Obese in Reception | | 9% | 9% | 10% | | 7% | 10% | | | 8% | Worsened |
| 11 % Obese in Year 6 | | 17% | 16% | 18% | | 15% | 23% | | | 15% | Worsened |
| 12 Teenage deliveries (age<19) in hospital per 1000 F15-17 | 36 | 12.6 | 14.7 | 21.5 | | 6.4 | | | | 13.1 | Worsened |
| 13 Emergency admission rate per 1000 for accidents (ages 0-17) | 288 | 9.4 | 9.3 | 10.9 | | 6.6 | | | | 9.2 | Worsened |
| 14 Admissions for self-harm rate per 1000 (ages 10-24) | 165 | 610.8 | 551.5 | 733.9 | | 242.5 | | 4421.2 | 61.7 | 451.2 | Worsened |
| 15 Emergency admissions rate per 1000 (ages 0-17) | 476 | 95 | 84 | 97 | | 64 | | 129 | 36 | 71 | Worsened |
| 16 Elective admissions rate per 1000 (aged 0-17) | 186 | 37 | 38 | 43 | | 27 | | 144 | 14 | 38 | Worsened |
| 17 First outpatient attendances rate per 1000 (aged 0-17) | 1,143 | 230 | 235 | 256 | | 201 | | 477 | 162 | 224 | Worsened |
| 18 % lone parent households | | 7% | 9% | 10% | | 7% | 11% | | | 6% | Worsened |
| 19 Foundation Stage Profile (FSP) % students NOT achieving a good level of development | | 50% | 33% | 50% | | 28% | | | | 60% | Worsened |
| 20 % of pupils with SEN | | 15% | 15% | 17% | | 12% | | | | 18% | Worsened |
| 21 Fixed Exclusions per 1000 pupils | | 41 | 46 | 52 | | 31 | | | | 37 | Worsened |
| 22 % Students NOT achieving 5 A*-C GCSEs inc Maths & English | | 50% | 41% | 50% | | 36% | | | | 58% | Worsened |
| 23 % of children in low-income families Child Poverty Unit 2014 | | 17% | 15% | 18% | | 13% | 20% | | | 15% | Worsened |
| 24 Children (0-17) currently in Care per 10,000** | 16 | 33 | 35 | 44 | | 19 | 60 | | | 28 | Worsened |
| 25 Children (0-17) subject to a Child Protection Plan per 10,000** | 11 | 22 | 30 | 61 | | 7 | 43 | | | 18 | Worsened |

* This indicator is arbitrarily drawn with a higher proportion shown on the left of the spine chart.

** These indicators are based on the previous configuration of practices in the old Localities.

Notes and Definitions

The values for the named Commissioning Locality (yellow circles) are compared with the Somerset averages (vertical line). Also shown (where data is available) is the England average (red diamonds). The range of Commissioning Locality values are shown by the grey bars. Better values are plotted to the right.

Some indicators are based on population weighted estimates and so do not have an actual number to show for Commissioning Localities or practices. Other practice data is missing where rates would have been based on very few events. England data is not available for locally calculated rates and some national data was not published because data was not complete.

- Proportion of population aged 0-14 as at April 2016 from Open Exeter GP registrations database.
- Income deprivation affecting Children Index 2015 Department for Communities and Local Government. Commissioning Locality value estimated using a weighted population average as at April 2016.
- Breast feeding initiation 2014/15. Estimated proportion of maternities where breastfeeding was initiated. NHS England.
- Breast feeding prevalence 2014/15. Estimated proportion of infants due checks who were reported as wholly or partially breastfed at 6-8 weeks. NHS England.
- Uptake of 5-in-1 primary vaccination by 1st birthday 2015/16 NHS England
- Uptake of 4-in-1 booster vaccination by 5th birthday 2015/16 NHS England
- Uptake of MMR primary vaccination by 2nd birthday 2015/16 NHS England
- Uptake of MMR booster vaccination by 5th birthday 2015/16 NHS England
- Uptake of vaccination by those aged 2, 3 and 4 during 2015/16 influenza season. NHS England.
- Proportion of children measured in Reception classes who had a Body Mass Index assessed as Obese for their age. 2010/11 - 2014/15 National Child Measurement Programme.
- Proportion of children measured in Year 6 classes who had a Body Mass Index assessed as Obese for their age. 2010/11 - 2014/15 National Child Measurement Programme.
- Number of deliveries in hospital to women aged less than 19, per 1000 females aged 15-17. April 2010 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- Emergency admissions for accidents per 1000 children aged 0-17. April 2010 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- Admissions for self-harm per 100,000 children and young people aged 10-24. April 2010 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- Emergency admissions for any cause per 1000 children aged 0-17. April 2015 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- Elective (day case or inpatient) admissions for any cause per 1000 children aged 0-17. April 2015 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- First outpatient attendances (a proxy for referral) for any cause per 1000 children aged 0-17. April 2015 to March 2016 Secondary Uses Services (SUS) hospital activity data.
- Proportion of households where there is a lone parent. Census 2011.
- Proportion of children not achieving a good level of development. Foundation Stage Profile (FSP). 2015 Somerset County Council.
- Proportion of children with Special Educational Needs. 2015 Somerset County Council.
- Number of Fixed exclusions from school per 1000 pupils. 2015 Somerset County Council.
- Proportion of children NOT achieving 5 A*-C GCSEs including Maths and English. 2015 Somerset County Council.
- Proportion of children in low-income families. Child Poverty Unit 2014.
- Children age 0-17 who are currently in Care and known to be living or placed in Somerset per 10,000. As at June 2016. Somerset County Council.
- Children age 0-17 who are subject to a Child Protection Plan and known to be living or placed in Somerset per 10,000. As at June 2016. Somerset County Council.

The profile contains information on the following lifestyle aspects of the local population:

- Obesity as assessed by the National Child Measurement Programme
- Obesity in adults
- Smoking prevalence
- Smoking cessation statistics
- Alcohol standardised admissions and mortality
- Drug misuse standardised admissions and mortality

Childhood obesity affects the ability to study, take part in physical activity and social activities. The National Child Measurement Programme (NCMP) measures the weight and height of children in reception class (aged 4 to 5 years) and year 6 (aged 10 to 11 years) to assess overweight children and obesity levels within primary schools. The measurement process is overseen by trained healthcare professionals in schools. The county council is responsible for the organisation of the local programme. Children's heights and weights are measured and used to calculate a Body Mass Index (BMI) centile based on the UK90 reference population. For population monitoring purposes, children above 85th centile are classified as overweight and above 95th centile as obese. Rates across England and Somerset show more children falling into the overweight and obese categories than might be expected. Population weighted averages of the rates in the areas where children live is used to estimate the rates in each Commissioning Locality.

Following the measurement programme parents are sent a feedback letter informing them of their child's weight and weight category (underweight, healthy weight, overweight and very overweight). The letters include links to web-based advice and support, contact numbers for local health teams and details of Somerset's children and families weight management programme. Copies of the feedback letters are also sent to GP practices.

Adult obesity is assessed based on raw BMI, kg/m². This data is taken from practice clinical records via the MIQUEST tool and looks at people measured as obese in the previous 15 months and so might not include all obese patients.

Smoking is the lifestyle feature which has the largest impact on individual health. Smoking ascertainment and prevalence are presented along with a range of statistics which highlight how well the smoking cessation process is operating. Somerset in general has lower rates of smoking than England but, despite recent improvements, still has much higher than average rates of smoking during pregnancy as assessed at time of delivery. More detail on local evidence based interventions to improve smokers' health is given on page 47.

Alcohol related admissions are more of a problem in Somerset than England as a whole and young people in particular show increased needs. There is increasing recognition of the damage that chronic excessive social drinking can cause. Although drug misuse affects a small proportion of the population than other lifestyle behaviours, the scale of impact can be far greater and it causes a disproportionate amount of human misery and takes a disproportionate amount of funding. Somerset Drug and Alcohol Service website provides a range of resources for professionals and the public to support healthier behaviour change <http://www.somersetdap.org.uk/>





West Somerset

Obesity in schools

National Child Measurement Programme (NCMP) Reception and Year 6 measurements 2011/12 - 2015/16

| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|-------------------------------|------------------------|----------|---------|--|
| Reception: Overweight | 13% | 14% | 13% | |
| Reception: Obese | 9% | 9% | 10% | |
| Reception: Overweight + Obese | 22% | 22% | 23% | |
| Year 6: Overweight | 14% | 14% | 14% | |
| Year 6: Obese | 17% | 16% | 23% | |
| Year 6: Overweight + Obese | 30% | 30% | 38% | |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show higher relative levels of overweight and obesity and so worse performance and values to the right show lower rates and so better performance.



West Somerset

Obesity in adults (rate per 1000)


Age/sex standardised prevalence rates using Quality Outcome Framework crude prevalence, Exeter system population downloads and Somerset wide age/sex specific rates from MIQUEST queries run by the CSU.

The Quality Outcome Framework indicators are reported annually through the Health and Social Care Information Centre website.

2016

| Condition | Observed in Commissioning Locality | Expected in Commissioning Locality (based on Somerset rates) | Commissioning Locality rate | Somerset rate | England rate | Range of Practice values low / median / high |
|---------------|------------------------------------|--|-----------------------------|---------------|--------------|--|
| Obesity (16+) | 3,307 | 2,778 | 105 | 88 | 104 | 33 / 81 / 235 |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show higher relative levels of overweight and obesity and so worse performance and values to the right show lower rates and so better performance.




Significantly worse than county average for:

Obesity (16+)

The Commissioning Locality has the worst value in the county for:

Obesity (16+)



Smoking prevalence (%)


Smokers 16+ among those with a known smoking status in the last 15 months from MIQUEST queries on practice clinical systems run by the CSU.

The value given is the number smoking of those of known smoking status in the last 15 months. This estimate is likely to be an overestimate as those who smoke are more likely to visit their GP and are more likely to have their smoking status recorded.


Mar-16

| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|--|------------------------|----------|---------|--|
| Number with known smoking status (16+) | 19,798 | 299,032 | | |
| Number smoking (16+) | 2,610 | 38,234 | | |
| Population (16+) | 29,429 | 473,432 | | |
| Proportion of population with known smoking status | 67% | 63% | | 49% / 63% / 74% |
| Estimated prevalence rate | 13.2% | 12.8% | | 4% / 12% / 29% |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show lower ascertainment or higher smoking rates and so worse performance.



The Commissioning Locality has the best value in the county for:
Proportion of population with known smoking status





Stop smoking services provision 2015/16

Smoking cessation services database - data is from a new provider

| | Commissioning Locality | Somerset | England | Range of Practice values low / median / high |
|---|------------------------|----------|---------|--|
| Number setting a quit date | 136 | 2,434 | | |
| Number going through a smoking cessation service per 1000 population per year | 4.3 | 4.4 | | 1 / 4 / 10 |
| Number quit | 79 | 1,184 | | |
| 4 week quit rate | 58% | 49% | | 8% / 48% / 100% |
| 4 week CO confirmed quit rate | 40% | 37% | | 0% / 35% / 100% |
| 4 week quit rate (45-64) | 42% | 45% | | |
| Smokers going through cessation per year per 1000 estimated smokers | 35 | 40 | | 5 / 38 / 99 |

The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. For consistency, values to the left show worse performance and to the right better performance which may be higher or lower absolute values.



Smoking evidence based interventions

1. **GPs to deliver brief or very brief interventions to identified smokers**, offering referral to those who express an interest in quitting, and offering help in the future for those who not ready to quit in the short term. VBI should be based on the NCSCT/BMJ Learning Module. Qualifies for 1 hour CPD/CME.

http://learning.bmj.com/learning/module-intro/advice-smoking.html?moduleid=10032720&locale=en_GB

Note that brief advice, alone, is significantly more effective than many standard medical treatments as shown in the table below, emphasising the importance of delivering this intervention well.

| Intervention | Outcome | NNT |
|--|---|------|
| Statins | Prevent one death over five years | 107 |
| Antihypertensive therapy | Prevent one stroke, MI, death over one year | 700 |
| Cervical cancer screening | Prevent one death over ten years | 1140 |
| GP brief advice to stop smoking (five minutes) | Prevent one premature death | 80 |

2. **Referral of smokers to stop smoking services.** Smokers who express an interest in quitting in the near future should be referred to the Smokefreelife Somerset service for support <http://www.smokefreelivesomerset.co.uk/>. In addition to group and 1-1 support, this service now includes telephone and video support. Smokers attending a NHS stop smoking service are up to 5 times more likely to quit than people quitting cold turkey or using OTC NRT. GPs should not normally prescribe stop smoking meds without behavioural support. If a smoker does not wish to be referred to a stop smoking practitioner, then it may be appropriate to provide a prescription, but other forms of support should be offered such as those available from the NHS Smokefree website, such as text, app and email. Repeat prescriptions should not be offered, with the patient referred to Smokefreelife Somerset to discuss options. There is a strong evidence base for the effectiveness of the text support service. Patients wishing to quit using electronic cigarettes/vaporisers will need to supply their own device, but are otherwise fully able to access Smokefreelife Somerset services.
3. **Practice records should be used to identify chronic disease patient groups for intervention.** In particular, patients with COPD who are continuing smokers should be a priority. Stop smoking groups specifically for COPD patients have been successful elsewhere, and would be worth trying in communities where there is sufficient interest. Smokefreelife Somerset should be contacted regarding facilitating such groups.
4. **Support for those likely to be hospitalised to quit smoking.** For patients likely to be admitted to hospital, GPs should ensure that patients are aware that hospitals are non-smoking sites (buildings and grounds) and that they will NOT be permitted to smoke anywhere on the site. They should then be offered support, either to quit prior to admission via Smokefreelife Somerset, or if unwilling to do so, should be prescribed NRT (preferably 2 products, patch plus a faster acting product such as lozenge) to enable temporary abstinence while in hospital, which they should start using a few days before admission.
5. **Engagement with the Stop Smoking Service provider.** Commissioning Localities and practices should work closely with Smokefreelife Somerset to ensure that stop smoking services are available to patients, and that referral processes are effective. The service has a GP and Pharmacy Co-ordinator, Sarah Westlake, who can be contacted via 01823 765006,

For the most part support is already available to practices for all the above from Smokefreelife Somerset. Public health can offer additional support if the Commissioning Localities wish to address any of the above at the larger geographic level. If you would like to work up any proposals please contact Stewart Brock, sbrock@somerset.gov.uk

Standardised admission and mortality rates for alcohol related conditions and drug misuse per 100,000 population per year

SUS Inpatient activity files April 2011 to March 2016 ONS Primary Care Mortality Database 2011-15. Population files July 2011-July 2015. Admissions indirectly standardised rate by age/sex and Deaths indirectly standardised rate by age/sex.


Alcohol: The North West Public Health Observatory has produced a list of conditions that they consider could have an alcohol related component. Each condition is allocated an Attributable Fraction, which is dependent on age and sex, that describes what proportion of admissions from that cause could be attributable to alcohol. These rates have been increasing year on year throughout the country. A very small percentage of admissions were excluded because of unknown age or gender.

Alcohol specific conditions are a subset of alcohol related conditions that can directly be attributable to alcohol.

Drug related harm: is represented according to the number of admissions which are judged as being related to drugs or deaths related to drug misuse. The National Treatment Agency for Substance Misuse guidance, Undertaking Needs Assessment – Drug Treatment (2009) recommended the following ICD10 codes: F10-F16, F18-F19, X42, X62 or Y12 in either the main diagnosis field or a secondary diagnosis field to identify substance misuse. **F10** relates to alcohol and is the major contributing code to substance misuse it is **excluded** in the following drug misuse indicators. A small number of admissions were excluded because of unknown age or gender.

| | Commissioning Locality rate | Somerset rate | England (average annual value 2010/11 to 2014/15) | Range of Practice values low / median / high |
|---|-----------------------------|---------------|---|--|
| Alcohol related admissions | 1,833 | 1,894 | 2,056 | 1,326 / 1,810 / 3,950 |
| Alcohol specific admissions | 495 | 445 | | 105 / 404 / 1,680 |
| Admissions for Alcoholic liver disease (K70) | 103 | 84 | | |
| Admissions for Alcoholic cirrhosis of liver (K703) | 59 | 44 | | |
| People admitted for Alcohol related conditions | 839 | 852 | | 636 / 824 / 1,853 |
| People admitted for Alcohol specific conditions | 271 | 248 | | 100 / 222 / 895 |
| People admitted for Alcoholic liver disease (K70) | 39 | 31 | | |
| People admitted for Alcoholic cirrhosis of liver (K703) | 27 | 16 | | |
| Alcohol related deaths | 3 | 6 | | |
| Drug misuse related admissions | 140 | 137 | | 20 / 109 / 488 |
| Drug misuse deaths | 3 | 3 | | |
| Alcohol specific OR drug misuse deaths | 40 | 31 | | |





The bar chart shows how the Commissioning Locality compares to other Commissioning Localities. Each vertical line on the yellow bar shows the position of a Commissioning Locality. Your Commissioning Locality is highlighted by the red diamond. Values to the left show higher rates of harm and so worse performance.



The Commissioning Locality has the worst value in the county for:

People admitted for Alcoholic liver disease (K70) People admitted for Alcoholic cirrhosis of liver (K703) Alcohol specific OR drug misuse deaths

Alcohol and drug misuse



The profile contains information on aspects relating to hospitalisation:

- Admission rates for specific conditions of interest: Self-harm and Mental and Behavioural disorders, falls, accidents to children, teenage deliveries
- Emergency admission rates
- Emergency admission rates for specific conditions of interest: CHD , CVD, COPD, Asthma and Diabetes
- Elective admissions rates
- Out patient attendances rates

This data enables scrutiny at a more local level of causes for emergency admissions. Emergency admissions – that is, admissions that are not predicted and happen at short notice because of perceived clinical need (NHS Connecting for Health 2010) – represent around 65 per cent of hospital bed days in England. In 2012-13, over a quarter of all patients attending major A&E departments were admitted, up from 19 per cent in 2003-04. This resulted in 5.3m emergency admissions at a cost of £12.5bn.

Whilst many admissions may be unavoidable, avoiding unnecessary emergency hospital admissions is a major concern for the NHS, not only because of the high and rising costs of emergency admission compared with other forms of care, but also because of the disruption it causes to elective health care – most notably inpatient waiting lists – and to the individuals admitted (Audit Commission 2009). It is recognised that a lack of alignment between hospitals and community and local services in the hours they are open compromises efforts to avoid out-of- hours hospital admissions and prolongs the length of stay of inpatients.

Rates of emergency admissions may indicate poorer patient management which has led up to a crisis point and thus indicate where improvements in primary care could be targeted to good effect. It is recommended that clinical commissioning groups to use data on variations in emergency admissions from ACSCs by constituent practices to understand variations in the quality of general practice as one of the causes. A recent NAO report (<http://www.nao.org.uk/wp-content/uploads/2013/10/10288-001-Executive-Summary.pdf>) estimates that 20% of admissions could be managed effectively in the community. Ambulatory care sensitive conditions (ACSCs) have been defined as chronic conditions for which it is possible to prevent acute exacerbations and reduce the need for hospital admission through active management, such as vaccination; better self-management, disease management or case management; or lifestyle interventions. Examples include congestive heart failure, diabetes, asthma, angina, epilepsy and hypertension. Focus on these conditions is one of the key proposed interventions to reduce overall emergency admissions. A 2012 Kings Fund briefing suggested that emergency admissions for ACSCs could be reduced by between 8 and 18 per cent resulting in savings of between £96 million and £238 million per year. Influenza, pneumonia, COPD, congestive heart failure, dehydration and gastroenteritis account for more than half (53 per cent) of the cost of emergency ACSCs admissions.

| The 19 ambulatory care-sensitive conditions - NHS Institute for Innovation and Improvement | |
|--|-------------------------------------|
| Vaccine-preventable | Acute |
| 1. Influenza and pneumonia | 11. Dehydration and gastroenteritis |
| 2. Other vaccine-preventable conditions | 12. Pyelonephritis |
| | 13. Perforated/bleeding ulcer |
| Chronic | 14. Cellulitis |
| 3. Asthma | 15. Pelvic inflammatory disease |
| 4. Congestive heart failure | 16. Ear, nose and throat infections |
| 5. Diabetes complications | 17. Dental conditions |
| 6. Chronic obstructive pulmonary disease (COPD) | 18. Convulsions and epilepsy |
| 7. Angina | 19. Gangrene |
| 8. Iron-deficiency anaemia | |
| 9. Hypertension | |
| 10. Nutritional deficiencies | |

For Somerset other indications are of relevance and interest such as rates of admissions for falls, especially rates in older people given the local population structure. Somerset has significantly higher rates of hospital admissions for self-harm than the rest of England. Guidance from NICE on common mental health disorders in primary care has been released at <http://pathways.nice.org.uk/pathways/common-mental-health-disorders-in-primary-care/common-mental-health-disorders-in-primary-care-overview>.

West Somerset

Standardised admission rates for self-harm and Mental and Behavioural disorders per 100,000 population per year




SUS Inpatient activity files April 2010 to March 2016 and ONS Population files July 2011-July 2015. Indirectly standardised rates by age/sex.

Admissions to hospital with any diagnosis of self-harm or self-injury of undetermined intent.

Admissions with a primary diagnosis in the Mental and Behavioural Disorders chapter of the International Classification of Diseases.

A very small percentage of self-harm admissions and mental and behavioural disorder admissions were excluded because of unknown age or gender.

| | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|---|-----------------------------|---------------|---------|--|
| Self-harm admissions | 206 | 221 | | 76 / 200 / 736 |
| Self-harm admissions for those aged 10-24 | 611 | 551 | | 62 / 513 / 4.421 |
| Mental and behavioural disorders admissions | 257 | 258 | | 105 / 235 / 829 |




West Somerset

Standardised emergency admission rates for Falls per 1000 population aged 65 and over and for accidents per 1000 population aged <18



SUS Inpatient activity files April 2013 to March 2016 and ONS Population files July 2013-July 2015. Indirectly standardised rates by age/sex.

Admissions to hospital with any diagnosis indicating a fall, for those aged 65 and over and any diagnosis indicating an Accident for those aged <18. No admissions were excluded because of unknown gender.

| | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|---------------|-----------------------------|---------------|---------|--|
| Falls 65+ | 26 | 31 | | 14 / 30 / 60 |
| Accidents <18 | 9.4 | 9.3 | | 4.2 / 9.2 / 14.7 |



Significantly better than county average for:
Falls 65+



Falls and Bone Health evidence based interventions

Falls are a major cause of injury, disability and mortality among older people, however there are many opportunities to reduce the risk of falls. It is also essential we support individuals to build and maintain optimum bone strength across the life course, assisting with maintenance of bone density into older age. During 2015/16 in Somerset approximately 4170 people aged 65 and over were admitted to hospital as a result of a fall and with our increasingly ageing population these numbers are set to continue to rise, we know that over 70% of falls go unreported placing an older person at risk of a more serious fall.

Hip fractures are costly both to the NHS and individuals. In Somerset we estimate each hip fracture costs the NHS £39,500 (2016) with additional substantial community hospital and social care costs, with 1430 hip fractures during 2015/16 this amounts to over £56.5 million for the NHS and £3 million for social care. In addition research has shown that up to 30% of people die within a year following a hip fracture and a further 20% have reduced capacity to live independently.

Alongside emphasis on falls reduction, diagnosing and treating osteoporosis is essential as 40-50% of women over 50 will experience a fragility fracture and 13-22% of men. Research also shows up to 50% of people who fracture a hip have sustained a previous fracture and a missed opportunity of diagnosis. In Somerset it is predicted 15,966 (12.6%) women have osteoporosis without a prior fracture and 14,949 (11.8%) women have osteoporosis with clinically apparent osteoporotic fragility fractures (2015).

Over the past few years national initiatives have been introduced and Somerset now has Bone Health and Falls pathways to provide comprehensive information and guidance for health professionals and other organisations. These are available on the Somerset Pathway Navigator App. The Bone Health pathway has sections on Healthy living for stronger bones; Who is at risk of osteoporosis; After fracture and End of life.

1) Primary diagnosis of osteoporosis

It is imperative we identify those with osteoporosis who are at risk of fragility fracture early on. Visit Somerset Bone Health pathway 'Who is at risk of osteoporosis' section via Navigator, this is supported by NICE Clinical Guidance CG 146 (2012) <http://publications.nice.org.uk/osteoporosis-assessing-the-risk-of-fragility-fracture-cg146>. The pathway suggests considering assessment in all women over 65 and men over 75, or younger if specific risk factors are present (including some health conditions/diseases – secondary causes, family history of osteoporosis, use of glucocorticoids, low BMI, smoking and more than 3 units of alcohol daily), using FRAX www.shef.ac.uk/FRAX/tool.jsp?country=1 to assess fracture risk, and a DXA scan where indicated.

2) Interventions including medication

When osteoporosis is diagnosed appropriate medication should be prescribed, as per NICE technology appraisals TA160 (2011) and TA204 (2010). Adherence to treatment regimens should be regularly reviewed. These guidelines also highlight the importance of a balanced diet with adequate levels of calcium and vitamin D and if people are not receiving an adequate amount of these nutrients through their diet prescribe vitamin D and calcium supplements. Further guidelines (2016) recommend adults in at risk groups take a vitamin D supplement of 10mcg per day particularly in the winter months. Safe exposure to sunlight is extremely important for the absorption of vitamin D and should be recommended in line with guidance.

3) Physical function and health issues

The Somerset Falls Pathway and Falls Risk Assessment Tool in Somerset (FRATIS) (visit Somerset Pathway Navigator App) assist in identifying specific risk factors and is used as a trigger for appropriate onward referral. This asks:

- Is there a history of any fall in the previous year?
- Is the patient/client on four or more medications per day?
- Does the patient/client have a diagnosis of stroke or Parkinson's Disease?
- Does the patient/client report any problems with his/her balance?
- Is the patient/client unable to rise from a chair of knee height (without assisting themselves)?
- It also includes a safe at home hazard assessment.

A positive response to three or more of the questions would indicate increased risk and a need for referral or other intervention.

4) Importance of physical activity

Evidence highlights that exercise programmes focusing on improving balance targeted at older high risk fallers, particularly those who have a history of falling, or a fear of falling are most effective in reducing falls and moderate injuries. A Cochrane review (2011) considered the evidence of exercise to improve bone health in postmenopausal women and found less BMD loss at the spine and neck of femur for those who were most active. Exercise also improves balance and maintains muscle strength. There are a range of exercise opportunities for individuals ranging from specialist Balance and Safety classes at Community Hospitals to community based Otago classes and modified tai chi and exercise for improving balance. The Zing website has details of all activity classes www.zingsomerset.co.uk.

5) Psychological aspects

It is widely recognised that psychological function has an impact on falls, this can be linked to specific cognitive problems but also to anxiety and a fear of falling, this can lead to avoidance of activity, further muscle and balance deterioration and reduced quality of life. It is important to break the cycle of anxiety; rehabilitation services can assist with confidence and increased mobility to help reduce the risk of social isolation and further deterioration.

6) Secondary prevention




Of vital importance to reduce hip fractures and other fragility fractures is diagnosis, management, and treatment of osteoporosis (see Somerset Bone Health Pathway in the Pathway Navigator App). Fracture Liaison Services (FLS) are now in place across Somerset and will identify all fractures in people over 50 and provide interventions, signposting, recommendations to GPs, and follow up where appropriate for a year after the fracture. Around fifty percent of people who fracture their hip have previously had a fragility fracture of some sort. This first, often relatively minor fracture (in terms of effect on morbidity and mortality) should be considered a warning sign of impending further fracture risk and trigger a comprehensive bone health and falls risk assessment. GPs are asked to action recommendations relating to prescribing and, after the first year on treatment, monitor patient medication adherence and suitability: seeking advice from the Bone Health Pathway; Falls Pathway; and/or FLS staff as appropriate.

Teenage deliveries in hospital

SUS Inpatient activity files April 2010 to March 2016. GP population files July 2010-July 2015. Rate per 1000 females aged 15-17 per year.

Numbers of deliveries in hospital where the age of the mother is less than 19 at delivery. This will not include home deliveries.

| | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|--------------------------|-----------------------------|---------------|---------|--|
| April 2010 to March 2016 | 12.6 | 14.7 | | |
| April 2010 to March 2013 | 16.1 | 18.2 | | |
| April 2013 to March 2016 | 8.8 | 11.0 | | |



West Somerset


Standardised emergency admission rate to hospital per 1000 population (ICD grouping)

SUS Inpatient activity files 2015/16. Population file July 2015. Emergency admissions. Standardised by age and sex.

A very small percentage of admissions are not included in the table below because the patient was of unknown age or sex.

| ICD groupings | Observed | Expected (based on Somerset rates) | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|---|--------------|------------------------------------|-----------------------------|---------------|---------|--|
| Diseases of the circulatory system | 456 | 469 | 10.6 | 10.9 | | 7.4 / 10.5 / 23.9 |
| Diseases of the respiratory system | 525 | 524 | 13.7 | 13.6 | | 6.5 / 13.3 / 23.0 |
| Diseases of the digestive system | 356 | 348 | 9.4 | 9.2 | | 4.0 / 9.2 / 13.5 |
| Diseases of musculoskeletal system and connective tissue | 165 | 195 | 4.1 | 4.9 | | 2.6 / 4.7 / 10.4 |
| Symptoms, signs and abnormal clinical and laboratory findings | 925 | 898 | 24.6 | 23.9 | | 13.6 / 23.1 / 51.7 |
| Injury, poisoning and other external causes | 635 | 641 | 16.8 | 16.9 | | 10.7 / 16.8 / 28.9 |
| Diabetes (any mention of diabetes for the admission) | 598 | 642 | 13.9 | 14.9 | | 6.2 / 14.1 / 29.6 |
| Falls (any mention of a fall for the admission) | 350 | 408 | 8.1 | 9.4 | | 5.2 / 9.3 / 14.8 |
| All | 4,267 | 4,270 | 112.1 | 112.2 | | 79 / 111 / 190 |
| All aged 75+ | 1,835 | 1,837 | 369.6 | 370.1 | | 265 / 372 / 596 |
| All aged <18 | 476 | 422 | 95.1 | 84.2 | | 36 / 82 / 129 |

Emergency admissions to hospital



Significantly worse than county average for:

All aged <18

Significantly better than county average for:

Falls (any mention of a fall for the admission)

West Somerset


Standardised emergency admission rate to hospital per 1000 population (CHD, CVD, COPD, Asthma, Diabetes)

SUS Inpatient activity files 2010/11 to 2015/16. Population file July 2010-July 2015. Emergency admissions. Standardised by age and sex.

A very small percentage of admissions are not included because of unknown age or sex.

| ICD groupings | Observed | Commissioning Locality rate | Somerset rate | England | Range of Practice values low / median / high |
|--|----------|-----------------------------|---------------|---------|--|
| Coronary heart disease | 558 | 2.0 | 2.3 | | 1.5 / 2.2 / 3.9 |
| Cerebrovascular disease | 626 | 2.2 | 1.9 | | 1.3 / 1.9 / 4.2 |
| COPD | 449 | 1.6 | 1.9 | | 0.8 / 1.8 / 7.6 |
| Asthma | 132 | 0.7 | 0.8 | | 0.2 / 0.7 / 1.5 |
| Diabetes (as the primary cause of admission) | 178 | 0.8 | 0.9 | | 0.2 / 0.9 / 2.4 |

Emergency admissions to hospital








Significantly worse than county average for:
Cerebrovascular disease

The Commissioning Locality has the worst value in the county for:
Cerebrovascular disease

Significantly better than county average for:
Coronary heart disease COPD Diabetes (as the primary cause of admission)

The Commissioning Locality has the best value in the county for:
COPD






Standardised elective admission rate to hospital per 1000 population (ICD grouping)

SUS Inpatient activity files 2015/16. Population file July 2015. Elective and Day cases. Standardised by age and sex.

A very small percentage of admissions are not included in the table below because the patient was of unknown age or sex.

| ICD groupings | Observed | Expected (based on Somerset rates) | Commissioning Locality rate | Somerset rate | England rate | Range of Practice values low / median / high |
|--|--------------|------------------------------------|-----------------------------|---------------|--------------|--|
| Cancer | 1,588 | 1,408 | 38.3 | 34.0 | | 15.7 / 34.3 / 61.0 |
| In situ and benign neoplasms and neoplasms of unknown or uncertain behaviour | 266 | 295 | 6.7 | 7.5 | | 3.4 / 6.8 / 16.2 |
| Diseases of eye and adnexa | 629 | 587 | 14.1 | 13.2 | | 6.0 / 13.5 / 19.8 |
| Diseases of the circulatory system | 356 | 347 | 8.8 | 8.5 | | 3.7 / 8.5 / 16.3 |
| Diseases of the digestive system | 894 | 955 | 24.2 | 25.8 | | 17.9 / 25.7 / 33.5 |
| Diseases of musculoskeletal system and connective tissue | 572 | 699 | 14.8 | 18.1 | | 12.0 / 18.1 / 25.8 |
| Symptoms, signs and abnormal clinical and laboratory findings | 356 | 383 | 9.3 | 10.0 | | 5.2 / 9.7 / 15.1 |
| Factors influencing health status and contact with health services | 406 | 311 | 10.6 | 8.1 | | 3.8 / 7.7 / 13.5 |
| Diabetes (any mention of diabetes for the admission) | 651 | 699 | 15.3 | 16.4 | | 9.2 / 16.0 / 27.7 |
| All | 6,233 | 6,273 | 159.8 | 160.8 | | 128 / 163 / 193 |
| All aged 75+ | 1,740 | 1,732 | 352.2 | 350.5 | | 204 / 351 / 507 |
| All aged <18 | 186 | 186 | 37.5 | 37.6 | | 14 / 37 / 144 |

Elective (inpatient and day cases) admissions to hospital



Significantly worse than county average for:
 Cancer Factors influencing health status and contact with health services

The Commissioning Locality has the worst value in the county for:
 Cancer

Significantly better than county average for:
 Diseases of musculoskeletal system and connective tissue

The Commissioning Locality has the best value in the county for:
 Diseases of musculoskeletal system and connective tissue


Standardised first outpatient attendance rate per 1000 population (proxy for referral)

SUS Outpatient activity files 2015/16. Population file July 2015. Standardised by age and sex.

Treatment function code is used rather than Specialty. The treatment function code reflects what the patient is treated for rather than the main specialty of the consultant under whose care the patient is. Not all treatment functions are listed. Some activity is only shown for Commissioning Localities. A small percentage of first attendances is not included in the table below because the patient was of unknown age or sex. Appropriate age and sex populations are used as the base for the rates.

| Treatment Function | Observed | Expected (based on Somerset rates) | Commissioning Locality rate | Somerset rate | England rate | Range of Practice values low / median / high |
|-----------------------|---------------|------------------------------------|-----------------------------|---------------|--------------|--|
| Trauma & Orthopaedics | 1,622 | 1,762 | 45.9 | 49.8 | | 40.5 / 49.7 / 63.7 |
| Ophthalmology | 1,438 | 1,463 | 35.4 | 36.0 | | 26.4 / 35.4 / 49.3 |
| Cardiology | 511 | 944 | 12.6 | 23.3 | | 8.0 / 17.6 / 85.0 |
| ENT | 735 | 842 | 20.8 | 23.8 | | 14.7 / 23.7 / 37.0 |
| Anaesthetics | 1,118 | 700 | 29.8 | 18.6 | | 0.2 / 20.2 / 34.9 |
| Respiratory medicine | 377 | 365 | 9.5 | 9.2 | | 3.2 / 9.2 / 19.7 |
| Gynaecology | 409 | 548 | 25.6 | 34.3 | | 17.1 / 31.0 / 71.3 |
| Urology | 590 | 655 | 14.9 | 16.6 | | 6.3 / 15.7 / 25.1 |
| Dermatology | 385 | 520 | 10.0 | 13.5 | | 6.4 / 12.7 / 23.7 |
| Physiotherapy | 434 | 539 | 12.0 | 14.9 | | 0.8 / 9.4 / 64.0 |
| Obstetrics | 235 | 342 | 49.7 | 72.4 | | 5.0 / 56.9 / 225.1 |
| Breast Surgery | 480 | 468 | 14.0 | 13.6 | | 8.7 / 13.4 / 18.4 |
| Paediatrics (<18) | 288 | 278 | 57.7 | 55.8 | | 29.9 / 53.8 / 94.8 |
| General Surgery | 263 | 326 | 7.1 | 8.7 | | 4.0 / 8.0 / 21.4 |
| Vascular Surgery | 408 | 381 | 9.7 | 9.0 | | 2.0 / 8.5 / 17.2 |
| Gastroenterology | 219 | 272 | 5.8 | 7.2 | | 2.7 / 6.6 / 15.0 |
| Clinical Oncology | 313 | 270 | 7.5 | 6.5 | | 1.6 / 6.4 / 13.8 |
| Colorectal Surgery | 301 | 294 | 7.7 | 7.5 | | 2.2 / 7.4 / 11.9 |
| All | 13,452 | 14,465 | 369.6 | 397.4 | | 308 / 389 / 524 |
| All 75+ | 3,330 | 2,956 | 866.9 | 769.6 | | 134 / 778 / 1361 |
| All <18 | 1,143 | 1,168 | 229.9 | 234.9 | | 162 / 234 / 307 |

First Outpatient attendances



Significantly worse than county average for:

Anaesthetics Clinical Oncology All 75+

Significantly better than county average for:

Trauma & Orthopaedics Cardiology ENT Gynaecology Urology Dermatology Physiotherapy Obstetrics General Surgery Gastroenterology All

The Commissioning Locality has the best value in the county for:

Cardiology ENT Dermatology Gastroenterology

The aim of the Medicines Optimisation Key Therapeutic Topics Comparators (formerly QIPP Prescribing Comparators) is to support organisations and prescribers to review the appropriateness of current prescribing, revise prescribing where appropriate and monitor implementation. The comparators are not intended to be used as targets or performance tables but rather highlight variation and support local discussion and decisions regarding MO KTT. More detail on the prescribing measures is available at <http://www.hscic.gov.uk/prescribing/measures> and the following data was taken from the website <https://apps.nhsbsa.nhs.uk/infosystems/welcome>:

ASTRO-PU weightings

ASTRO-PU stands for Age, Sex and Temporary Resident Originated Prescribing Units. This weighting is designed to weight individual practice or organisation populations for age and sex to allow for better comparison of prescribing patterns. These figures are based on the cost or volume of prescribing across all therapeutics, and these weightings should be used only when considering all prescribing. The number of temporary residents attending practices is no longer captured or included in funding allocations. The cost based weightings are standardised (based on a male child under 4 years being 1) as they are used in national resource allocation formulae. The item based weightings are not standardised, as this more clearly shows relative use across different demographic groups.

MO KTT Prescribing indicators

NHS Business Authority Information Portal (via CCG)


For the 4 indicators at the top of the table and on the left of the graph a higher rate is better. For the other 9 indicators a lower rate is better.

2015/16

| Indicator | Numerator | Denominator | Commissioning Locality rate | Somerset rate | England rate | Range of Practice values low / median / high |
|--|-----------|-------------|-----------------------------|---------------|--------------|--|
| 3 day courses of antibiotics: ADQ/item | 18,719 | 3,195 | 5.9 | 5.9 | 5.9 | 4.6 / 5.9 / 8.0 |
| Antidepressants: First choice % items | 17,715 | 26,542 | 66.7 | 62.7 | 69.1 | 29 / 66 / 86 |
| Blood glucose lowering drugs | 15,624 | 18,703 | 83.5 | 79.7 | 80.4 | 69 / 81 / 91 |
| NSAIDs: Ibuprofen & Naproxen % Items | 8,744 | 10,552 | 82.9 | 78.4 | 78.6 | 56 / 80 / 91 |
| Antibacterial Items/STAR PU | 19,295 | 21,436 | 0.9 | 1.0 | 1.1 | 0.7 / 1.0 / 3.2 |
| Antidepressants ADQ/STAR PU | 913,533 | 801,907 | 1.1 | 1.3 | 1.4 | 0.9 / 1.3 / 1.9 |
| Co-Amoxiclav, Cephalosporins and Quinolones: % Items | 982 | 19,247 | 5.1 | 6.9 | 9.7 | 2.9 / 6.4 / 14.3 |
| Dosulepin: % items | 526 | 26,542 | 2.0 | 1.4 | 2.1 | 0.0 / 1.2 / 5.7 |
| Hypnotics ADQ/STAR PU (ADQ based) | 101,460 | 123,896 | 0.8 | 0.8 | 1.0 | 0.3 / 0.8 / 2.3 |
| Long/Intermediate Insulin Analogues | 827 | 1,544 | 53.6 | 67.7 | 78.7 | 39 / 68 / 89 |
| Minocycline ADQ/1000 Patients | 458 | 29 | 15.5 | 12.2 | 40.3 | 0 / 0 / 265 |
| NSAIDs: ADQ/STAR PU | 306,501 | 54,269 | 5.6 | 5.9 | 5.6 | 3.0 / 5.6 / 15.2 |
| Other Lipid Modifying Drugs: % items | 882 | 48,054 | 1.8 | 1.9 | 2.2 | 0.3 / 1.6 / 4.3 |

Orange highlighting indicates an improvement over the previous year.

MO KTT prescribing data - latest year





The Commissioning Locality has the worst value in the county for:

Dosulepin: % items

The Commissioning Locality has the best value in the county for:

Blood glucose lowering drugs NSAIDs: Ibuprofen & Naproxen % Items Antidepressants ADQ/STAR PU Co-Amoxiclav, Cephalosporins and Quinolones: % Items Long/Intermediate Insulin Analogues



2014/15

| Indicator | Numerator | Denominator | Commissioning Locality rate | Somerset rate | England rate | Range of Practice values low / median / high |
|--|-----------|-------------|-----------------------------|---------------|--------------|--|
| 3 day courses of antibiotics: ADQ/item | 19,094 | 3,165 | 6.0 | 6.1 | 6.0 | 4.7 / 6.1 / 7.7 |
| Antidepressants: First choice % items | 14,991 | 25,616 | 58.5 | 57.1 | 63.7 | 33 / 59 / 80 |
| Blood glucose lowering drugs | 15,734 | 18,668 | 84.3 | 81.7 | 82.5 | 69 / 83 / 90 |
| NSAIDs: Ibuprofen & Naproxen % Items | 8,858 | 10,781 | 82.2 | 77.7 | 76.2 | 59 / 79 / 91 |
| Antibacterial Items/STAR PU | 20,532 | 21,462 | 1.0 | 1.1 | 1.2 | 0.8 / 1.0 / 3.5 |
| Antidepressants ADQ/STAR PU | 860,965 | 803,852 | 1.1 | 1.2 | 1.3 | 0.8 / 1.2 / 1.8 |
| Co-Amoxiclav, Cephalosporins and Quinolones: % Items | 1,483 | 20,450 | 7.3 | 9.0 | 10.7 | 3.9 / 8.4 / 17.0 |
| Dosulepin: % items | 562 | 25,616 | 2.2 | 1.8 | 2.6 | 0.0 / 1.5 / 6.7 |
| Hypnotics ADQ/STAR PU (ADQ based) | 110,012 | 123,718 | 0.9 | 0.9 | 1.1 | 0.2 / 0.8 / 2.6 |
| Long/Intermediate Insulin Analogues | 791 | 1,452 | 54.5 | 70.8 | 80.0 | 41 / 71 / 94 |
| Minocycline ADQ/1000 Patients | 392 | 33 | 11.8 | 20.9 | 50.9 | 0 / 1 / 468 |
| NSAIDs: ADQ/STAR PU | 317,052 | 54,127 | 5.9 | 6.0 | 5.9 | 3.0 / 6.0 / 14.7 |
| Other Lipid Modifying Drugs: % items | 886 | 48,254 | 1.8 | 2.0 | 2.4 | 0.5 / 1.7 / 4.9 |

West Somerset

ASTRO-PUs at April 2016 **281,239**


Age/Sex ASTRO-PU pyramid
 Solid line represents Somerset as a whole, dotted line represents England.
 Data from Exeter system download of GP registered patients.

ASTRO-PUs here are units based on the population at a Commissioning Locality and are used to help with prescribing budgets. Weighting of the population is higher for the sections of the population expected to need more prescribing spend.

West Somerset

| Age Group | Males | Females |
|-----------|--------|---------|
| 00-04 | 672 | 495 |
| 05-14 | 1,516 | 1,336 |
| 15-24 | 2,232 | 3,106 |
| 25-34 | 3,022 | 4,404 |
| 35-44 | 4,017 | 5,740 |
| 45-54 | 10,496 | 13,597 |
| 55-64 | 22,420 | 24,883 |
| 65-74 | 43,948 | 41,789 |
| 75-84 | 32,578 | 34,188 |
| 85+ | 11,542 | 19,259 |

The ASTRO-PU distribution reflects the Commissioning Locality population profile.



Somerset

| Age Group | Males | Females |
|-----------|---------|---------|
| 00-04 | 14,717 | 11,297 |
| 05-14 | 34,574 | 30,024 |
| 15-24 | 44,559 | 59,592 |
| 25-34 | 55,917 | 86,806 |
| 35-44 | 90,419 | 127,808 |
| 45-54 | 202,164 | 252,119 |
| 55-64 | 332,893 | 360,787 |
| 65-74 | 550,665 | 527,365 |
| 75-84 | 390,962 | 404,244 |
| 85+ | 139,682 | 219,577 |

